

BRX-TH-504
BOW-PH-125
hep-th/0206094

pp-wave limits and orientifolds

Stephen G. Naculich^a, Howard J. Schnitzer^{1,b}, and Niclas Wyllard^{2,b}

^a*Department of Physics
Bowdoin College, Brunswick, ME 04011*

^b*Martin Fisher School of Physics
Brandeis University, Waltham, MA 02454*

Abstract

We study the pp-wave limits of various elliptic models with orientifold planes and D7-branes, as well as the pp-wave limit of an orientifold of $adS_5 \times T^{11}$. Many of the limits contain both open and closed strings. We also present pp-wave limits of theories which give rise to a compact null direction and contain open strings. Maps between the string theory states and gauge theory operators are proposed.

¹Research supported in part by the DOE under grant DE-FG02-92ER40706.

² Research supported by the DOE under grant DE-FG02-92ER40706.
naculich@bowdoin.edu; schnitzer,wyllard@brandeis.edu

Contents

1	Introduction	2
2	Gauge theories on branes in orientifold backgrounds	3
3	Penrose limits of orientifolds of $adS_5 \times S^5$	4
4	$\mathcal{N} = 4$ $Sp(2N)$ and $SO(N)$ gauge theories	6
5	$\mathcal{N} = 2$ $Sp(2N)$ with $\square \oplus 4\square$	7
6	$\mathcal{N} = 2$ $SU(N)$ with $2\square \oplus 4\square$	9
7	The $\mathcal{N} = 2$ $SU \times SU \times \dots \times SU \times SU$ orientifold	16
8	$\mathcal{N} = 2$ $Sp(2N) \times Sp(2N)$ with $(\square, \square) \oplus 2(1, \square) \oplus 2(\square, 1)$	19
9	The $\mathcal{N} = 2$ $Sp \times SU \times \dots \times SU \times Sp$ orientifold	24
10	$\mathcal{N} = 1$ $Sp(2N) \times Sp(2N)$ with $2(\square, \square) \oplus 4(1, \square) \oplus 4(\square, 1)$	26
11	Summary	29

1 Introduction

Recently a particular limiting version of the adS/CFT correspondence [1] between string theory on $adS_5 \times S^5$ and the $\mathcal{N} = 4$ supersymmetric gauge theory was discovered [2] (for earlier work see refs. [3, 4, 5]). On the string theory side the limiting metric is the metric experienced by an observer moving very fast along a particular null geodesic of $adS_5 \times S^5$, viz., a parallel plane (pp) wave metric [6]. The limiting background also contains a non-zero RR five-form. We will refer to this limit interchangeably as a Penrose or pp-wave limit. Type IIB string theory in the pp-wave background is exactly solvable [5, 7]. Moreover, the spectrum of string theory states in this background corresponds to the subset of the single trace operators in the gauge theory with large conformal dimension Δ and R-charge J , but with $H \equiv \Delta - J$ finite. The operators in the superconformal $\mathcal{N} = 4$ $SU(N)$ gauge theory that correspond to the string theory states in the Penrose limit of $adS_5 \times S^5$ were identified in ref. [2].

In this paper we study examples of the limiting procedure introduced in ref. [2] for a class of models containing orientifold planes (most of the models also contain D7-branes) and therefore potentially having an open string sector.

Sections 2 and 3 contain background material relevant for our later discussions, with section 3 exhibiting the two different classes of Penrose limits encountered in this paper. In section 4 we consider the pp-wave limit of the simplest orientifolded model, $adS_5 \times \mathbb{RP}^5$ and its dual $\mathcal{N} = 4$ SO/Sp gauge theory. In section 5 we discuss some aspects of the simplest orientifold of $adS_5 \times S^5$ containing D7-branes, whose dual is the $\mathcal{N} = 2$ $Sp(2N)$ gauge theory

with $\square \oplus 4\square$ matter hypermultiplets. The pp-wave limit containing an open string sector was studied in ref. [8]; we focus on another Penrose limit of the same theory, which does not contain open strings. (Other aspects of open strings and D-branes in the pp-wave background have been discussed in refs. [9].)

In sections 6 and 8 we discuss the pp-wave limits of theories whose IIB descriptions contain (before taking the limit) an O7-plane, four D7-branes (plus mirror branes), and a $\mathbb{Z}_2$ orbifold; specifically, $\mathcal{N} = 2$ $SU(N)$ with $2\square \oplus 4\square$ matter hypermultiplets, and $\mathcal{N} = 2$ $Sp(2N) \times Sp(2N)$ with $(\square, \square) \oplus 2(1, \square) \oplus 2(\square, 1)$ matter hypermultiplets. The adS/CFT correspondence for these elliptic models was discussed in detail in refs. [10, 11]. Both theories possess two different Penrose limits, one of which allows for an open string sector. Moreover, in these models there are several distinct species of open strings, due to the presence of the orbifold projection.

In sections 7 and 9 we discuss two infinite classes of elliptic models [12] for which the models considered in sections 6 and 8 are the first members. The type IIB descriptions of these two classes of models contain an O7-plane, four D7-branes (plus mirror branes), and a $\mathbb{Z}_{2N_2}$ orbifold, and the dual gauge theories have gauge groups SU^{N_2} and $Sp \times SU^{N_2-1} \times Sp$, respectively. We show that these theories possess a scaling limit, similar to the one recently considered in refs. [13, 14], giving rise to a compact null direction. The theories we consider contain both open and closed string sectors.

In section 10 we discuss the pp-wave limit for the correspondence between type IIB string theory on $adS_5 \times T^{11}/\mathbb{Z}_2^{\text{ori}}$ and the $\mathcal{N} = 1$ $Sp(2N) \times Sp(2N)$ theory with $2(\square, \square) \oplus 4(1, \square) \oplus 4(\square, 1)$ matter chiral multiplets. By studying this example we are able to test the ideas of ref. [8] in a more complicated setting. Section 11 summarizes the results of this paper.

2 Gauge theories on branes in orientifold backgrounds

The four-dimensional $\mathcal{N} = 4$, $\mathcal{N} = 2$, and $\mathcal{N} = 1$ supersymmetric gauge theories considered in this paper arise as the worldvolume theories on D3-branes in various orientifolded type IIB backgrounds. The $\mathcal{N} = 4$ and $\mathcal{N} = 2$ theories arise from D3-branes in a flat background modded out by an orientifold group G . We parameterize the flat six-dimensional space transverse to the D3-brane worldvolume by

$$\begin{aligned} z_1 = x_6 + ix_7 &= r \cos \theta_1 \cos \theta_2 e^{i\varphi_1}, & \theta_{1,2} &\in [0, \tfrac{1}{2}\pi], \\ z_2 = x_8 + ix_9 &= r \cos \theta_1 \sin \theta_2 e^{i\varphi_2}, & \varphi_{1,2,3} &\in [0, 2\pi], \\ z_3 = x_4 + ix_5 &= r \sin \theta_1 e^{i\varphi_3}. \end{aligned} \quad (2.1)$$

The coordinates z_i are subject to various identifications that depend on the form of G .

In the large- N limit ($\sim$ large number of D3-branes), the D3-branes significantly modify the background. The near-horizon limit of the resulting background is the space $adS_5 \times S^5$, whose metric is given by

$$ds^2 = R^2 (ds_{adS_5}^2 + d\tilde{\Omega}_5^2), \quad (2.2)$$

$$\begin{aligned} ds_{adS_5}^2 &= d\rho^2 - \cosh^2 \rho dt^2 + \sinh^2 \rho d\Omega_3^2, \\ d\tilde{\Omega}_5^2 &= d\theta_1^2 + \cos^2 \theta_1 (d\theta_2^2 + \cos^2 \theta_2 d\varphi_1^2 + \sin^2 \theta_2 d\varphi_2^2) + \sin^2 \theta_1 d\varphi_3^2. \end{aligned} \quad (2.3)$$

Modding out by the orientifold group G results in identifications among the coordinates, specified in each case below.

The isometry group of S^5 is $\text{SO}(6)$, which is to be identified with the $\text{SU}(4)$ R -symmetry group of the $\mathcal{N} = 4$ theories. For the $\mathcal{N} = 2$ theories considered in this paper, G reduces the isometry group to $[\text{SO}(4) \times \text{SO}(2)]/G$, where $\text{SO}(4) \simeq \text{SU}(2)_L \times \text{SU}(2)_R$ is the rotation group on $\mathbb{C}^2 \sim \{z_1, z_2\}$, and $\text{SO}(2) \simeq \text{U}(1)_R$ is the rotation group on $\mathbb{C} \sim \{z_3\}$. More precisely, $\text{SU}(2)_L$ and $\text{SU}(2)_R$ act on the left and right of the matrix

$$\begin{pmatrix} z_1 & -\bar{z}_2 \\ z_2 & \bar{z}_1 \end{pmatrix}. \quad (2.4)$$

The diagonal generators $J_{L,R}^3$ of $\text{SU}(2)_{L,R}$ and the generator J_R of $\text{U}(1)_R$, act as

$$J_R^3 + J_L^3 = -i\partial_{\varphi_1}, \quad J_R^3 - J_L^3 = -i\partial_{\varphi_2}, \quad J_R = -i\partial_{\varphi_3}. \quad (2.5)$$

$\text{SU}(2)_R \times \text{U}(1)_R/G$ is the R -symmetry group of the $\mathcal{N} = 2$ field theories, whereas $\text{SU}(2)_L$ is a global symmetry group. For some of the models we consider $\text{SU}(2)_L$ is further reduced to $\text{U}(1)_L$. For these models J_L^3 is identified with the $\text{U}(1)_L$ generator. To facilitate comparison to the results in ref. [11], we note that $J_R^3 = \frac{1}{2}q_R$.

In this paper, we also consider the $\mathcal{N} = 1$ $\text{Sp}(2N) \times \text{Sp}(2N)$ field theory on a stack of D3-branes in the background of an orientifold of the conifold. The orientifold group and the gauge theory on the D3-branes are discussed in more detail in section 10.

3 Penrose limits of orientifolds of $adS_5 \times S^5$

All geodesics of S^5 are equivalent, so there is only one Penrose limit of $adS_5 \times S^5$, when the geodesic lies partially in the S^5 directions. (If the geodesic lies completely inside adS_5 , the Penrose limit gives flat space [15].) For the orientifolded theories considered in this paper, however, Penrose limits along different geodesics can give rise to different string theories. In this section, we identify two particular geodesics and the associated Penrose limits.

Penrose limit A

Consider the geodesic parameterized by φ_3 and contained in the subspace $\rho = 0$, $\theta_1 = \frac{1}{2}\pi$. Define

$$x^+ = \frac{1}{2}(t + \varphi_3), \quad x^- = \frac{1}{2}R^2(t - \varphi_3), \quad r = R\rho, \quad y = R(\frac{1}{2}\pi - \theta_1). \quad (3.6)$$

In the limit $R \rightarrow \infty$ (keeping $x^\pm$, r , and y finite), the metric goes to

$$ds^2 = -4dx^+dx^- - (r^2 + y^2)(dx^+)^2 + dr^2 + r^2d\tilde{\Omega}_3^2 + dy^2 + y^2d\tilde{\Omega}_3^2, \quad (3.7)$$

where

$$d\tilde{\Omega}_3^2 = d\theta_2^2 + \cos^2\theta_2 d\varphi_1^2 + \sin^2\theta_2 d\varphi_2^2. \quad (3.8)$$

The space transverse to the null geodesic is $\mathbb{R}^4 \times \tilde{\mathbb{R}}^4$. Parameterizing $\tilde{\mathbb{R}}^4 = \mathbb{C}^2$ by

$$w_1 = y \cos\theta_2 e^{i\varphi_1}, \quad w_2 = y \sin\theta_2 e^{i\varphi_2}, \quad (3.9)$$

we can rewrite the metric as

$$ds^2 = -4dx^+dx^- - (r^2 + |w_1|^2 + |w_2|^2)(dx^+)^2 + dr^2 + r^2d\Omega_3^2 + |dw_1|^2 + |dw_2|^2. \quad (3.10)$$

The p^- lightcone momentum is given by $2p^- = H_A = i\partial_{x^+} = i(\partial_t + \partial_{\varphi_3})$, which is identified in the dual gauge theory with [2] $H_A = \Delta - J_A$, where Δ is the conformal dimension and J_A is a global symmetry generator. From (2.5) we see that $J_A = J_R$. Hence, in Penrose limit A, $H_A = \Delta - J_R$.

Penrose limit B

Consider the geodesic parameterized by φ_1 lying along $\rho = 0$ and $\theta_1 = \theta_2 = 0$. Define

$$x^+ = \frac{1}{2}(t + \varphi_1), \quad x^- = \frac{1}{2}R^2(t - \varphi_1), \quad r = R\rho, \quad y_2 = R\theta_2, \quad y_3 = R\theta_1. \quad (3.11)$$

In the limit $R \rightarrow \infty$ (keeping $x^\pm$, r , y_2 , and y_3 finite), the metric goes to

$$ds^2 = -4dx^+dx^- - (r^2 + y_2^2 + y_3^2)(dx^+)^2 + dr^2 + r^2d\Omega_3^2 + dy_2^2 + y_2^2d\varphi_2^2 + dy_3^2 + y_3^2d\varphi_3^2. \quad (3.12)$$

The space transverse to the null geodesic is $\mathbb{R}^4 \times \tilde{\mathbb{R}}^4$. Parameterizing $\tilde{\mathbb{R}}^4 = \mathbb{C}^2$ by

$$w_2 = y_2e^{i\varphi_2}, \quad w_3 = y_3e^{i\varphi_3}, \quad (3.13)$$

the metric becomes

$$ds^2 = -4dx^+dx^- - (r^2 + |w_2|^2 + |w_3|^2)(dx^+)^2 + dr^2 + r^2d\Omega_3^2 + |dw_2|^2 + |dw_3|^2. \quad (3.14)$$

From $2p^- = H_B = i(\partial_t + \partial_{\varphi_1})$ we see that $J_B = -i\partial_{\varphi_1}$, which using (2.5), implies that $J_B = J_R^3 + J_L^3$. Hence, in Penrose limit B, $H_B = \Delta - J_R^3 - J_L^3$.

The Penrose limits discussed here are limits of the full supergravity/string theory background (as discussed in ref. [3]). The non-trivial RR five-form field strength is also non-zero in the limit (and is responsible for giving masses to the lightcone fermions). As observed in ref. [2], string theory in the pp-wave background is dual to a certain subset of gauge theory operators.

The metrics in the two Penrose limits (3.10) and (3.14) are identical. However, the orientifold group G imposes different identifications on the coordinates in the two cases, and thus can lead to distinct string theories. These distinct Penrose limits therefore correspond to different subsets of operators in the gauge theory on the D3 branes in the orientifolded background.

Finally, we note that the limit along the geodesic parameterized by φ_2 is equivalent to the limit involving φ_1 (Penrose limit B). On the gauge theory side, the only difference is a trivial sign redefinition of J_L^3 .

In section 10 we discuss the Penrose limit of an orientifold of $adS_5 \times T^{11}$. More details about the orientifold group and the Penrose limit are given in that section.

4 $\mathcal{N} = 4$ $\text{Sp}(2N)$ and $\text{SO}(N)$ gauge theories

Let us start by considering the $\mathcal{N} = 4$ $\text{SO}(N)$ and $\text{Sp}(2N)$ theories. In $\mathcal{N} = 1$ language, these theories contain a vector multiplet, and three chiral multiplets ϕ_{ia}^b . The scalars in the chiral multiplets, organized into $SU(2)_R$ multiplets, and their quantum numbers are shown in the table below.

CFT field	$\text{Sp}(2N)$ or $\text{SO}(N)$	J_R	J_R^3	J_L^3	Δ	H_A	H_B
ϕ_3	adjoint	1	0	0	1	0	1
$\phi_3^\dagger$	adjoint	-1	0	0	1	2	1
$(\phi_1, \phi_2^\dagger)$	adjoint	0	$(\frac{1}{2}, -\frac{1}{2})$	$\frac{1}{2}$	1	(1,1)	(0,1)
$(\phi_2, \phi_1^\dagger)$	adjoint	0	$(\frac{1}{2}, -\frac{1}{2})$	$-\frac{1}{2}$	1	(1,1)	(1,2)

Table 1: $SU(2)_R$ multiplets of scalar fields in $\mathcal{N} = 4$ $\text{Sp}(2N)$ or $\text{SO}(N)$.

The $\mathcal{N} = 4$ $\text{SO}(N)$ and $\text{Sp}(2N)$ gauge theories arise as the worldvolume theories on a stack of D3-branes parallel to an O3-plane in a flat type IIB background. The generator $\Omega(-1)^{F_L} R_{456789}$ of the orientifold group $\mathbb{Z}_2^{\text{ori}}$ acts on the $\mathbb{C}^3$ transverse to the D3-branes as

$$z^1 \rightarrow -z^1, \quad z^2 \rightarrow -z^2, \quad z^3 \rightarrow -z^3, \quad (4.1)$$

where R_{456789} denotes reflection in the directions 4, $\dots$, 9, Ω is the worldsheet parity operation, and $(-1)^{F_L}$ changes the sign of the left-movers in the Ramond sector.

In the near-horizon limit, the geometry becomes $adS_5 \times S^5 / \mathbb{Z}_2^{\text{ori}}$ where the geometric part of $\mathbb{Z}_2^{\text{ori}}$ acts as

$$\varphi_1 \rightarrow \varphi_1 + \pi, \quad \varphi_2 \rightarrow \varphi_2 + \pi, \quad \varphi_3 \rightarrow \varphi_3 + \pi, \quad (4.2)$$

on the coordinates of S^5 (2.3), which yields $adS_5 \times \mathbb{RP}^5$. The adS/CFT correspondence between string theory on this background and the above SO/Sp gauge theories was developed in ref. [16]. The Penrose limit of these theories was briefly studied in ref. [17]; below we give some further details.

Penrose limit A

The action (4.2) of $\mathbb{Z}_2^{\text{ori}}$ on the coordinates in Penrose limit A is an inversion of $\tilde{\mathbb{R}}^4$: $w_1 \rightarrow -w_1$, $w_2 \rightarrow -w_2$, accompanied by a shift $\varphi_3 \rightarrow \varphi_3 + \pi$ halfway around the geodesic (see section 3 for details about the notation). This $\mathbb{Z}_2^{\text{ori}}$ action has no fixed points on S^5 , and therefore no fixed points in the Penrose limit (as we will see in later sections, this fact is related to the absence of an open string sector). We now discuss the operators in the gauge theory that correspond to the states in the closed string theory.

The only candidate for the operator corresponding to the vacuum state of the closed string theory in the pp-wave background (3.10) is¹

$$\text{tr} \left[\phi_3^{J_A} \right], \quad (4.3)$$

¹Throughout this paper we will not keep track of the overall normalizations of the operators.

which has $\Delta = J_A$ and therefore $H_A = 0$. The effect of modding out by $\mathbb{Z}_2^{\text{ori}}$ is to halve the periodicity of the geodesic circle, and thus to allow only even integer values of J_A for the vacuum string states. Thus from the string theory perspective one expects the vacuum to correspond to

$$\text{tr} \left[\phi_3^{2n} \right]. \quad (4.4)$$

From the gauge theory perspective, the exponent must be even because the adjoint fields satisfy $\phi_i^T = J\phi_i J$ ($\phi_i^T = -\phi_i$) in the Sp theory (SO theory), so that $\text{tr}(\phi_3^{2n+1})$ vanishes identically.

Consider string states in which a zero-momentum oscillator $a_0^{\dagger\mu}$ in one of the directions of $\mathbb{R}^4$ acts on the light-cone vacuum. These states correspond to the insertion of $(D_\mu\phi_3)\phi_3$ into (4.4), yielding four operators

$$\text{tr} \left[(D_\mu\phi_3)\phi_3^{2n+1} \right], \quad (4.5)$$

which have $\Delta = 2n + 3$ and $J_A = 2n + 2$, and therefore $H_A = 1$.

Next, consider string states in which a zero-momentum oscillator $a_0^{\dagger i}$, $\bar{a}_0^{\dagger i}$ in one of the directions w_i , $\bar{w}_i$ ($i = 1, 2$) of $\tilde{\mathbb{R}}^4$ acts on the light-cone vacuum. The inversion $(w_1, w_2) \rightarrow (-w_1, -w_2)$ causes $a_0^{\dagger i} \rightarrow -a_0^{\dagger i}$, $\bar{a}_0^{\dagger i} \rightarrow -\bar{a}_0^{\dagger i}$, so for the string state to be invariant under $\mathbb{Z}_2^{\text{ori}}$, the value of J_A must be odd. These string states correspond to insertions of $\phi_1\phi_3$, $\phi_1^\dagger\phi_3$, $\phi_2\phi_3$, and $\phi_2^\dagger\phi_3$ into (4.4), yielding four additional $H_A = 1$ operators

$$\text{tr} \left[\phi_1\phi_3^{2n+1} \right], \quad \text{tr} \left[\phi_1^\dagger\phi_3^{2n+1} \right], \quad \text{tr} \left[\phi_2\phi_3^{2n+1} \right], \quad \text{tr} \left[\phi_2^\dagger\phi_3^{2n+1} \right]. \quad (4.6)$$

Again, from the gauge theory perspective, the exponents follow from the Sp (SO) constraints on the ϕ_i 's.

The eight operators (4.5) and (4.6) therefore represent the eight bosonic zero-momentum states in the string theory. The correspondence for the fermionic oscillators as well as for the non-zero-momentum oscillators can be constructed in complete analogy with the original construction [2]; we will not give the details here.

For the case of $adS_5 \times \mathbb{RP}^5$, the distinction between the Sp and SO theories was explained in ref. [16]. An interesting question is to understand how they are distinguished in the pp-wave limit.

Penrose limit B

The action (4.2) of $\mathbb{Z}_2^{\text{ori}}$ on the coordinates in Penrose limit B involves an inversion of the transverse $\mathbb{C}^2$: $w_2 \rightarrow -w_2$, $w_3 \rightarrow -w_3$, accompanied by a shift $\varphi_1 \rightarrow \varphi_1 + \pi$ halfway around the geodesic. This action is identical to the one in Penrose limit A after a trivial relabelling of the coordinates. The correspondence of string states to the gauge theory operators is the same as in Penrose limit A except with ϕ_1 and ϕ_3 interchanged.

5 $\mathcal{N} = 2$ Sp(2N) with $\square \oplus 4\square$

The $\mathcal{N} = 2$ Sp(2N) theory with one antisymmetric and four fundamental hypermultiplets contains an $\mathcal{N} = 2$ vector multiplet in the adjoint representation, which in $\mathcal{N} = 1$ language

consists of a vector multiplet and a chiral multiplet, ϕ_a^b , in the adjoint representation. The theory also contains an antisymmetric $\mathcal{N} = 2$ hypermultiplet, which in $\mathcal{N} = 1$ language consists of a chiral multiplet in the $\square$ representation of $\text{Sp}(2N)$, $A_{ab} = -A_{ba}$, and a chiral multiplet in the $\bar{\square}$ representation $\tilde{A}^{ab} = -\tilde{A}^{ba}$. The indices on the latter can be lowered using the symplectic unit J_{ab} , leaving us with two chiral multiplets, $A_{1ab} = A_{ab}$ and $A_{2ab} = J_{ac}J_{bd}\tilde{A}^{ab}$, in the $\square$ representation, transforming as a doublet of $\text{SU}(2)_L$. Finally, the theory contains four fundamental $\mathcal{N} = 2$ hypermultiplets, which in $\mathcal{N} = 1$ language consists of eight chiral multiplets Q_a^I in the $\square$ representation (after lowering the indices of the four chiral multiplets in the $\bar{\square}$ representation). The scalars in the chiral multiplets and their quantum numbers are shown in the table below.

CFT field	$\text{Sp}(2N)$	J_R	J_R^3	J_L^3	Δ	H_A	H_B
ϕ	adjoint	1	0	0	1	0	1
$\phi^\dagger$	adjoint	-1	0	0	1	2	1
$(A_1, (JA_2J)^\dagger)$	$\square$	0	$(\frac{1}{2}, -\frac{1}{2})$	$\frac{1}{2}$	1	(1,1)	(0,1)
$(A_2, (JA_1J)^\dagger)$	$\square$	0	$(\frac{1}{2}, -\frac{1}{2})$	$-\frac{1}{2}$	1	(1,1)	(1,2)
$(Q^I, Q^{I\dagger})$	$\square$	0	$(\frac{1}{2}, -\frac{1}{2})$	0	1	1	$(\frac{1}{2}, \frac{3}{2})$

Table 2: $\text{SU}(2)_R$ multiplets of scalar fields in $\mathcal{N} = 2$ $\text{Sp}(2N)$ with $\square \oplus 4\square$.

The above $\mathcal{N} = 2$ $\text{Sp}(2N)$ gauge theory arises [18] as the theory on a stack of D3-branes parallel to an O7-plane and four physical D7-branes in a flat background. The orientifold group $\mathbb{Z}_2^{\text{ori}} = \{\mathbb{1}, \Omega R_{45}(-1)^{F_L}\}$, where R_{45} reflects the 45 directions, acts on the space transverse to the D3-branes as $z^3 \rightarrow -z^3$. This action fixes the hyperplane $z_3 = 0$, which corresponds to the position of the O7-plane and the D7-branes.

In the near-horizon limit the background becomes $adS_5 \times S^5 / \mathbb{Z}_2^{\text{ori}}$, where $\mathbb{Z}_2^{\text{ori}}$ acts as $\varphi_3 \rightarrow \varphi_3 + \pi$ on the coordinates of S^5 . The fixed point set of this action is an S^3 located at $\theta_1 = 0$. The adS/CFT correspondence for this model was discussed in ref. [19].

Penrose limit A

In Penrose limit A, the generator of $\mathbb{Z}_2^{\text{ori}}$ simply produces a translation along the geodesic $\varphi_3 \rightarrow \varphi_3 + \pi$ (together with the action of $\Omega(-1)^F$) with no action on the transverse coordinates w_1 and w_2 (3.9). This limit therefore yields the maximally supersymmetric pp-wave background. However, $\mathbb{Z}_2^{\text{ori}}$ projects out half of the discrete values of J_A that would be allowed in the unprojected case.

Closed string sector

The $H_A = 0$ operators

$$\text{tr} [\phi^{2n}], \quad (5.1)$$

correspond to the closed string vacuum states, where the effect of modding out by $\varphi_3 \rightarrow \varphi_3 + \pi$ is to allow only even integer values of J_A . From the gauge theory perspective, the exponent

must be even because the adjoint field satisfies $\phi^T = J\phi J$, implying that $\text{tr}[\phi^{2n+1}]$ vanishes identically.

String states in which a zero-momentum oscillator $a_0^{\dagger\mu}$ in one of the directions of $\mathbb{R}^4$ acts on the light-cone vacuum correspond to the $H_A = 1$ gauge theory operators

$$\text{tr}[(D_\mu\phi)\phi^{2n+1}]. \quad (5.2)$$

String states in which a zero-momentum oscillator $a_0^{\dagger i}, \bar{a}_0^{\dagger i}$ in one of the directions of $\tilde{\mathbb{R}}^4$ acts on the light-cone vacuum correspond to the four $H_A = 1$ operators

$$\text{tr}[(A_i J)\phi^{2n}], \quad \text{tr}[(JA_i^\dagger)\phi^{2n}], \quad (i = 1, 2), \quad (5.3)$$

where we write $A_i J$ for $A_{iac}J^{cb}$ (and similarly for $JA_i^\dagger$). Using $(AJ)^T = -J(AJ)J$, it follows that the operator would vanish if the exponent of ϕ were odd.

Non-zero-momentum states can be constructed in complete analogy with the original construction [2].

The fixed point set (O7-plane) of $\mathbb{Z}_2^{\text{ori}}$ goes away in Penrose limit A, and so do the D7-branes, so there is no open string sector. The absence of an open string sector corresponds to the fact that the operator

$$Q^I J \phi^n Q^K \quad (5.4)$$

vanishes by using the F-term equations of the gauge theory. This implies that it is not a chiral primary operator, and therefore can not correspond to a vacuum state.

Penrose limit B

In Penrose limit B, the geometric action of $\mathbb{Z}_2^{\text{ori}}$, $\varphi_3 \rightarrow \varphi_3 + \pi$ produces a reflection of the transverse coordinate $w_3 \rightarrow -w_3$. The geodesic lies in the orientifold plane $w_3 = 0$, and the coincident D7-branes gives rise to an open string sector in addition to the closed string sector.

This limit of the theory was considered in detail by Berenstein et. al. [8], to which we refer the reader for a complete description of the closed and open string sectors and the corresponding gauge theory operators.

6 $\mathcal{N} = 2$ $\text{SU}(N)$ with $2\Box \oplus 4\Box$

The $\mathcal{N} = 2$ $\text{SU}(N)$ theory with two antisymmetric and four fundamental hypermultiplets contains an $\mathcal{N} = 2$ vector multiplet in the adjoint representation, which in $\mathcal{N} = 1$ language comprise a vector multiplet and a chiral multiplet, ϕ_a^b , in the adjoint representation. The theory also contains two antisymmetric $\mathcal{N} = 2$ hypermultiplets, which in $\mathcal{N} = 1$ language comprise a pair of chiral multiplets A_{iab} ($i = 1, 2$) in the $\Box$ representation and a pair of chiral multiplets $\tilde{A}_i^{ab}$ ($i = 1, 2$) in the $\bar{\Box}$ representation. Both (A_1, A_2) and $(\tilde{A}_1, \tilde{A}_2)$ form $\text{SU}(2)_L$ doublets. The conjugate fields $(A^{\dagger 1}, A^{\dagger 2})$ transform in the conjugate $\text{SU}(2)_L$ representation, but by lowering the indices with the ϵ tensor, so that $A^{\dagger 1} = A_2^\dagger$ and $A^{\dagger 2} = -A_1^\dagger$ we can form

an $SU(2)_L$ doublet $(A_1^\dagger, A_2^\dagger)$ from the conjugate fields (and similarly for $(\tilde{A}_1^\dagger, \tilde{A}_2^\dagger)$). Finally, the theory contains four fundamental $\mathcal{N} = 2$ hypermultiplets, which in $\mathcal{N} = 1$ language comprise four chiral multiplets Q_a^I in the $\square$ representation and four chiral multiplets $\tilde{Q}_I^a$ in the $\overline{\square}$ representation. The quantum numbers of the scalars in the above supermultiplets are shown in the table below.

CFT field	$SU(N)$	J_R	J_R^3	J_L^3	Δ	H_A	H_B
ϕ	adjoint	1	0	0	1	0	1
$\phi^\dagger$	adjoint	-1	0	0	1	2	1
$(A_1, \tilde{A}_1^\dagger)$	$\square$	0	$(\frac{1}{2}, -\frac{1}{2})$	$\frac{1}{2}$	1	(1,1)	(0,1)
$(A_2, \tilde{A}_2^\dagger)$	$\square$	0	$(\frac{1}{2}, -\frac{1}{2})$	$-\frac{1}{2}$	1	(1,1)	(1,2)
$(\tilde{A}_1, A_1^\dagger)$	$\overline{\square}$	0	$(\frac{1}{2}, -\frac{1}{2})$	$\frac{1}{2}$	1	(1,1)	(0,1)
$(\tilde{A}_2, A_2^\dagger)$	$\overline{\square}$	0	$(\frac{1}{2}, -\frac{1}{2})$	$-\frac{1}{2}$	1	(1,1)	(1,2)
$(Q^I, \tilde{Q}_I^\dagger)$	$\square$	0	$(\frac{1}{2}, -\frac{1}{2})$	0	1	(1,1)	$(\frac{1}{2}, \frac{3}{2})$
$(\tilde{Q}_I, Q^{I\dagger})$	$\overline{\square}$	0	$(\frac{1}{2}, -\frac{1}{2})$	0	1	(1,1)	$(\frac{1}{2}, \frac{3}{2})$

Table 3: $SU(2)_R$ multiplets of scalar fields in $\mathcal{N} = 2$ $SU(N)$ with $2\square \oplus 4\overline{\square}$.

The F-term equations for the model are

$$A_1\tilde{A}_2 - A_2\tilde{A}_1 + Q^I\tilde{Q}_I = 0, \quad \phi A_i + A_i\phi^T = 0, \quad \tilde{A}_i\phi + \phi^T\tilde{A}_i = 0, \quad \phi Q^I = 0 = \tilde{Q}_I\phi. \quad (6.1)$$

The $\mathcal{N} = 2$ $SU(N)$ gauge theory just described arises as the worldvolume theory on a stack of coincident D3-branes in a flat type IIB background modded out by the orientifold group

$$G = \mathbb{Z}_2^{\text{orb}} \times \mathbb{Z}_2^{\text{ori}}, \quad (6.2)$$

where $\mathbb{Z}_2^{\text{orb}} = \{\mathbb{1}, R_{6789}\}$ and $\mathbb{Z}_2^{\text{ori}} = \{\mathbb{1}, R_{45}\Omega(-1)^{F_L}\}$ act on the $\mathbb{C}^3$ transverse to the D3-branes as

$$z_3 \xrightarrow{\mathbb{Z}_2^{\text{ori}}} -z_3, \quad (z_1, z_2) \xrightarrow{\mathbb{Z}_2^{\text{orb}}} (-z_1, -z_2). \quad (6.3)$$

The fixed point set of $\mathbb{Z}_2^{\text{ori}}$ is the hyperplane $z_3 = 0$, which corresponds to the position of an O7-plane and four physical D7-branes, while the fixed point set of $\mathbb{Z}_2^{\text{orb}}$ is the six-dimensional hyperplane $z_1 = z_2 = 0$.

The fields of the $SU(N)$ theory discussed above can be obtained via a projection from those of the $\mathcal{N} = 4$ $SU(2N)$ theory², whose three adjoint $\mathcal{N} = 1$ chiral superfields are denoted Φ_i . The independent generators γ_θ and $\gamma_{\Omega'}$ of the orientifold group are realized as the $2N \times 2N$ matrices [12]

$$\gamma_\theta = i \begin{pmatrix} \mathbb{1}_N & 0 \\ 0 & -\mathbb{1}_N \end{pmatrix}, \quad \gamma_{\Omega'} = \begin{pmatrix} 0 & \mathbb{1}_N \\ -\mathbb{1}_N & 0 \end{pmatrix}, \quad (6.4)$$

²The projection really gives a $U(N)$ theory; however, since we are interested in the large- N limit the $U(1)$ factor is suppressed and can be ignored.

which induce the following projections on the fields of the gauge theory:

$$\begin{aligned}\Phi_{1,2} &= -\gamma_\theta \Phi_{1,2} \gamma_\theta^{-1}, & \Phi_3 &= \gamma_\theta \Phi_3 \gamma_\theta^{-1}, \\ \Phi_{1,2} &= \gamma_{\Omega'} \Phi_{1,2}^T \gamma_{\Omega'}^{-1}, & \Phi_3 &= -\gamma_{\Omega'} \Phi_3^T \gamma_{\Omega'}^{-1},\end{aligned}\tag{6.5}$$

resulting in Φ_i 's of the form

$$\Phi_1 = \begin{pmatrix} 0 & A_1 \\ -\tilde{A}^2 & 0 \end{pmatrix} = \begin{pmatrix} 0 & A_1 \\ \tilde{A}_1 & 0 \end{pmatrix}, \quad \Phi_2 = \begin{pmatrix} 0 & A_2 \\ \tilde{A}^1 & 0 \end{pmatrix} = \begin{pmatrix} 0 & A_2 \\ \tilde{A}_2 & 0 \end{pmatrix}, \quad \Phi_3 = \begin{pmatrix} \phi & 0 \\ 0 & -\phi^T \end{pmatrix}.\tag{6.6}$$

In the near-horizon (large- N) limit of the above orientifolded background one obtains $adS_5 \times S^5 / [\mathbb{Z}_2^{\text{orb}} \times \mathbb{Z}_2^{\text{ori}}]$, where $\mathbb{Z}_2^{\text{orb}}$ acts as $\varphi_1 \rightarrow \varphi_1 + \pi$, $\varphi_2 \rightarrow \varphi_2 + \pi$ and $\mathbb{Z}_2^{\text{ori}}$ acts as $\varphi_3 \rightarrow \varphi_3 + \pi$ on the coordinates of S^5 . String theory in this background is dual to the $SU(N)$ gauge theory discussed above (see ref. [11] for further details). Below we discuss the Penrose limits for this correspondence.

Penrose limit A

In the first Penrose limit, the generator of $\mathbb{Z}_2^{\text{ori}}$ simply produces a translation along the geodesic: $\varphi_3 \rightarrow \varphi_3 + \pi$ (together with the action of $\Omega(-1)^{F_L}$), with no action on the transverse coordinates w_1 and w_2 (3.9). There is no orientifold fixed plane and therefore no open string sector in this limit, but $\mathbb{Z}_2^{\text{ori}}$ projects out half of the discrete values of J_A that would be allowed in the unprojected case. The generator of $\mathbb{Z}_2^{\text{orb}}$ acts solely on the transverse coordinates as $w_1 \rightarrow -w_1$, $w_2 \rightarrow -w_2$. The geodesic lies in the orbifold fixed plane, resulting in a twisted sector of the string theory in this pp-wave background. First we describe the operators corresponding to the untwisted sector.

Untwisted sector

The identification of the gauge theory operators corresponding to the states of the dual string theory in the Penrose limit is facilitated by going over to the cover space, as in refs. [20, 13] (see also [21]). The ground state in the untwisted sector corresponds to $\text{tr}[\Phi_3^{J_A}]$. As a result of the orientifold projection (6.6), this is only nonvanishing for even J_A , yielding the $H_A = 0$ operator

$$\text{tr} [\phi^{2n}].\tag{6.7}$$

The $H_A = 1$ gauge theory operators $\text{tr}[(D_\mu \Phi_3) \Phi_3^{2n+1}]$, or equivalently using (6.6),

$$\text{tr} [(D_\mu \phi) \phi^{2n+1}],\tag{6.8}$$

correspond to string states in which a single zero-momentum oscillator $a_0^{\dagger\mu}$ in one of the directions of $\mathbb{R}^4$ acts on the light-cone vacuum.

Since the zero-momentum oscillators $a_0^{\dagger I}$ in the $\tilde{\mathbb{R}}^4$ directions (3.9) are odd under $\mathbb{Z}_2^{\text{orb}}$, an even number of $a_0^{\dagger I}$'s must act on the light-cone vacuum. The $H_A = 2$ gauge theory operators corresponding to the states in which two such oscillators act on the vacuum involve

the insertion of two $H_A = 1$ fields $\Upsilon_I = (\Phi_1, \Phi_1^\dagger, \Phi_2, \Phi_2^\dagger)$ into the ground state operator (the insertion of a single $H_A = 1$ operator into the ground state operator gives zero using (6.6)):

$$\sum_{m=0}^{J_A} \text{tr} \left[\Upsilon_I \Phi_3^m \Upsilon_K \Phi_3^{J_A-m} \right], \quad (6.9)$$

where we average over the relative position of the Υ 's [2, 22, 23]. The sum over m vanishes unless $J_A = 2n$. Using (6.6), together with

$$\Phi_1^\dagger = \begin{pmatrix} 0 & \tilde{A}_1^{\dagger 1} \\ A_1^{\dagger 1} & 0 \end{pmatrix} = \begin{pmatrix} 0 & \tilde{A}_2^{\dagger 1} \\ A_2^{\dagger 1} & 0 \end{pmatrix}, \quad \Phi_2^\dagger = \begin{pmatrix} 0 & \tilde{A}_1^{\dagger 2} \\ A_1^{\dagger 2} & 0 \end{pmatrix} = \begin{pmatrix} 0 & -\tilde{A}_1^{\dagger 1} \\ -A_1^{\dagger 1} & 0 \end{pmatrix}, \quad (6.10)$$

we obtain the set of ten operators

$$\begin{aligned} \sum_m (-)^m \text{tr} \left[\tilde{A}_{(i} \phi^m A_{j)} (\phi^T)^{2n-m} \right], \quad \sum_m (-)^m \text{tr} \left[A_{(i}^\dagger \phi^m \tilde{A}_{j)}^\dagger (\phi^T)^{2n-m} \right], \\ \sum_m (-)^m \text{tr} \left[A_i^\dagger \phi^m A_j (\phi^T)^{2n-m} + \tilde{A}_j \phi^m \tilde{A}_i^\dagger (\phi^T)^{2n-m} \right]. \end{aligned} \quad (6.11)$$

These correspond to the ten string theory states

$$\begin{aligned} a_0^{\dagger i} a_0^{\dagger j} |0, p^+\rangle, \quad \bar{a}_0^{\dagger i} \bar{a}_0^{\dagger j} |0, p^+\rangle, \\ \bar{a}_0^{\dagger i} a_0^{\dagger j} |0, p^+\rangle, \end{aligned} \quad (6.12)$$

where $a_0^{\dagger i}$ ($\bar{a}_0^{\dagger i}$) is the zero-momentum operator in the direction w_i ($\bar{w}_i$).

In limit A there are no open strings, since the D7-branes disappear. The gauge theory explanation of this fact is that the candidate open string vacuum $\tilde{Q}_I \phi^{2n} Q^K$ is ruled out since it is not a chiral primary operator, as can be seen from the fact that it vanishes by the F-term equation, $\phi Q^K = 0$.

Twisted sector

The ground state in the twisted sector of the closed string theory corresponds to the operator $\text{tr}[\gamma_\theta \Phi_3^{J_A}]$. The presence of the operator γ_θ in the trace requires J_A to be odd, giving

$$\text{tr} \left[\phi^{2n+1} \right]. \quad (6.13)$$

The $H_A = 1$ operators corresponding to the string theory oscillators in the $\mathbb{R}^4$ directions of the pp-wave metric are identified, in the twisted sector, with $\text{tr}[\gamma_\theta (D_\mu \Phi_3) \Phi_3^{2n+2}]$, or

$$\text{tr} \left[(D_\mu \phi) \phi^{2n+2} \right]. \quad (6.14)$$

The oscillators in the twisted sector are half-integer moded [24, 25, 20, 26], so there are no zero-momentum oscillators in this sector. Hence the operators that would have corresponded to such oscillators acting on the twisted-sector vacuum

$$\sum_m \text{tr} \left[\gamma_\theta \Upsilon_I \Phi_3^m \Upsilon_J \Phi_3^{2n-m} \right], \quad (6.15)$$

must not be chiral primary operators in the gauge theory. Consider for example

$$\sum_m (-)^m \text{tr} \left[\tilde{A}_{[i} \phi^m A_{j]} (\phi^T)^{2n-m} \right]. \quad (6.16)$$

By using the F-term equation $\tilde{A}_i \phi + \phi^T \tilde{A}_i = 0$, $\tilde{A}_i$ can be shifted through the ϕ^T 's. Then $A_{[j} \tilde{A}_{i]}$ can be replaced by Q 's, using $A_1 \tilde{A}_2 - A_2 \tilde{A}_1 + Q^I \tilde{Q}_I = 0$. Finally, using $\phi Q^I = 0$, the operator vanishes and is therefore not a chiral primary operator. The other operators in (6.15) are similarly expected to be ruled out.

Although there are no zero-momentum oscillators in the twisted sector, non-zero modes are present. The construction of the corresponding gauge theory operators proceeds in analogy with the discussions in refs. [24, 25, 20, 26].

Penrose limit B

In limit B, the geometric action of $\mathbb{Z}_2^{\text{ori}}$, $\varphi_3 \rightarrow \varphi_3 + \pi$, produces a reflection of the coordinate transverse to the O7-plane: $w_3 \rightarrow -w_3$, cf. (3.13). Thus the geodesic lies in the orientifold (O7) plane $w_3 = 0$, and the coincident D7-branes give rise to an open-string sector. The generator of $\mathbb{Z}_2^{\text{orb}}$ produces a shift $\varphi_1 \rightarrow \varphi_1 + \pi$ halfway around the geodesic, accompanied by the reflection $w_2 \rightarrow -w_2$. Since the $\mathbb{Z}_2^{\text{orb}}$ action has no fixed points, there is no twisted sector in this Penrose limit, however $\mathbb{Z}_2^{\text{orb}}$ projects out half of the discrete values of J_B that would be allowed in the unprojected case. The pp-wave background in this limit is the same as the one considered in ref. [8] except for the additional action of $\mathbb{Z}_2^{\text{orb}}$, which leads to different species of open string states, not encountered in ref. [8].

Closed string sector

The vacuum state of the closed string sector corresponds to the $H_B = 0$ operator $\text{tr}[\Phi_1^{J_B}]$, which, due to the $\mathbb{Z}_2^{\text{orb}}$ projection, is only nonvanishing when $J_B = 2n$,

$$\text{tr}[\Phi_1^{2n}] = \text{tr} \left[(A_1 \tilde{A}_1)^n \right]. \quad (6.17)$$

(On the gauge theory side the constraint on J_B is required because we need an equal number of $\square$ and $\bar{\square}$ fields to form a gauge invariant operator.) The absence of the twisted sector vacuum can be seen from the fact that $\text{tr}[\gamma_\theta \Phi_1^m] = 0$ for all values of m .

String states in which a zero-momentum oscillator $a_0^\dagger{}^\mu$ in one of the directions of $\mathbb{R}^4$ acts on the light-cone vacuum correspond to the $H_B = 1$ gauge theory operator

$$\text{tr}[(D_\mu \Phi_1) \Phi_1^{2n+1}] = \text{tr} \left[D_\mu (A_1 \tilde{A}_1) (A_1 \tilde{A}_1)^n \right]. \quad (6.18)$$

String states in which a zero-momentum oscillator $a_0^\dagger$ ($\bar{a}_0^\dagger$) in the w_2 ($\bar{w}_2$) direction of $\tilde{\mathbb{R}}^4$ acts on the light-cone vacuum corresponds to insertions of the $H_B = 1$ field $\Phi_2 \Phi_1$ ($\Phi_2^\dagger \Phi_1$) into the vacuum state operator:

$$\begin{aligned} \text{tr} \left[\Phi_2 \Phi_1^{2n+1} \right] &= \text{tr} \left[(A_2 \tilde{A}_1 + A_1 \tilde{A}_2) (A_1 \tilde{A}_1)^n \right], \\ \text{tr} \left[\Phi_2^\dagger \Phi_1^{2n+1} \right] &= \text{tr} \left[(\tilde{A}_1^\dagger \tilde{A}_1 + A_1 A_1^\dagger) (A_1 \tilde{A}_1)^n \right]. \end{aligned} \quad (6.19)$$

Note that an insertion of $(A_1\tilde{A}_2 - A_2\tilde{A}_1)$ can be replaced by $Q^I\tilde{Q}_I$ via the F-term equations, and thus turns the would-be $H_B = 1$ closed string state into the (subleading) open string vacuum (6.23) (this appears to be related to the closed/open string interaction). Since the operator vanishes modulo subleading terms by using the F-term equations it is not protected and is thus ruled out. There is no similar gauge theory argument which rules out the insertion of the $H_B = 1$ operator $A_1A_1^\dagger - \tilde{A}_1^\dagger\tilde{A}_1$. It would be interesting to better understand why this operator is ruled out³.

The zero-momentum oscillators in the w_3 , $\bar{w}_3$ directions of $\tilde{\mathbb{R}}^4$ correspond to insertions of ϕ and $\phi^\dagger$ into the vacuum state (6.17). These oscillators are odd under $\mathbb{Z}_2^{\text{ori}}$ and so there should be no operators corresponding to single oscillators acting on the vacuum. This is automatic in the cover space language since the gauge theory operators $\text{tr}[\Phi_3\Phi_1^{2n}]$ and $\text{tr}[\Phi_3^\dagger\Phi_1^{2n}]$ vanish identically. An independent argument can also be constructed. The insertion of a single ϕ is not possible since $\text{tr}[\phi(A_1\tilde{A}_1)^n] = \frac{1}{2}\text{tr}[(\phi A_1 + A_1\phi^T)\tilde{A}_1(A_1\tilde{A}_1)^{(n-1)}]$, which vanishes by the F-term equations. Thus the operator is not a chiral primary (nor is it a “near chiral primary”) and is consequently ruled out. We expect that the insertion of a single $\phi^\dagger$ is also ruled out as implied by the cover space result.

States with two oscillators in the w_3 direction acting on the ground state are not projected out and correspond to the insertion of a pair of Φ_3 ’s

$$\sum_{m=0}^{2n} \text{tr} [\Phi_3\Phi_1^m\Phi_3\Phi_1^{2n-m}], \quad (6.20)$$

which gives

$$\sum_{m \text{ even}} \text{tr} \left[\phi (A_1\tilde{A}_1)^{\frac{m}{2}} \phi (A_1\tilde{A}_1)^{n-\frac{m}{2}} \right] - \sum_{m \text{ odd}} \text{tr} \left[\phi (A_1\tilde{A}_1)^{\frac{m-1}{2}} \tilde{A}_1\phi^T\tilde{A}_1 (A_1\tilde{A}_1)^{n-\frac{m+1}{2}} \right]. \quad (6.21)$$

States where one (or both) of the zero-momentum oscillators are in the $\bar{w}_3$ direction are obtained from the above expressions by replacing one (or both) of the Φ_3 ’s by $\Phi_3^\dagger$.

Open string sector

Since the geodesic lies in the orientifold (O7) fixed-plane $w_3 = 0$ which is also the location of the D7-branes, there is an open-string sector. The gauge symmetry of the D7-branes/O7-plane, $\text{SO}(8)_F$, is broken to $\text{SU}(4)_F \times \text{U}(1)$ by the $\mathbb{Z}_2^{\text{orb}}$ projection. The ground state of the open string sector, which transforms in the adjoint **28** of $\text{SO}(8)_F$, decomposes into

$$\mathbf{28} \rightarrow \mathbf{15}_0 \oplus \mathbf{1}_0 \oplus \mathbf{6}_2 \oplus \bar{\mathbf{6}}_{-2}. \quad (6.22)$$

The $\mathbb{Z}_2^{\text{orb}}$ projection correlates the states in the $\mathbf{15}_0 \oplus \mathbf{1}_0$ representation with operators in odd-dimensional representations of $\text{SU}(2)_L$ [11]. These open string vacuum states should therefore correspond to gauge theory operators where the total number of A ’s and $\tilde{A}$ ’s is even. There is a natural candidate

$$\tilde{Q}_I(A_1\tilde{A}_1)^n Q^K, \quad (6.23)$$

³A similar comment applies to the $H_B = 1$ insertion, $(D_\mu A_1)\tilde{A}_1 - A_1(D_\mu\tilde{A}_1)$.

which indeed belongs to the correct $(\mathbf{15}_0 \oplus \mathbf{1}_0)$ $SU(4)_F \times U(1)$ representation. Furthermore, the operator (6.23) has $H_B = 1$, reflecting the non-vanishing zero point energy for the open string sector [8].

The $\mathbb{Z}_2^{\text{orb}}$ projection correlates the states in the $\mathbf{6}_2 \oplus \bar{\mathbf{6}}_{-2}$ representation with operators in even-dimensional representations of $SU(2)_L$ [11]. These open string vacuum states should therefore correspond to gauge theory operators where the total number of A 's and $\tilde{A}$'s is odd. The natural candidates are

$$Q^I \tilde{A}_1 (A_1 \tilde{A}_1)^n Q^K, \quad \tilde{Q}_{[I} (A_1 \tilde{A}_1)^n A_1 \tilde{Q}_{K]}. \quad (6.24)$$

In these expressions, the flavor indices on the Q 's are antisymmetrized because the operator sandwiched in between them is antisymmetric in gauge indices, which shows that the operators (6.24) belong to the correct $SU(4)$ representations. Furthermore, they have the correct $U(1)$ (quark number) charges.

Notice that in contrast to the model discussed in [8] there are two distinct types of open strings in this model. The first, corresponding to (6.23) are “meson” open strings with a quark at one end and an anti-quark at the other, whereas the other, corresponding to (6.24), have quark number ± 2 , with quarks (or anti-quarks) at both ends.

The open string state in which a zero-momentum oscillator in one of the $\mathbb{R}^4$ directions act on the open string vacua corresponds to the insertion of the operator $D_\mu(A_1 \tilde{A}_1)$ into the operators in (6.23) and (6.24).

Zero-momentum oscillators $a_0^\dagger$, $\bar{a}_0^\dagger$ in the w_2 , $\bar{w}_2$ directions of $\tilde{\mathbb{R}}^4$ acting on the meson string vacuum (6.23) correspond to the two $H_B = 2$ operators

$$\sum_m \tilde{Q}_I (A_1 \tilde{A}_1)^m (A_1 \tilde{A}_2 + A_2 \tilde{A}_1) (A_1 \tilde{A}_1)^{n-m} Q^K, \quad (6.25)$$

$$\sum_m \tilde{Q}_I (A_1 \tilde{A}_1)^m (\tilde{A}_1^\dagger \tilde{A}_1 + A_1 A_1^\dagger) (A_1 \tilde{A}_1)^{n-m} Q^K, \quad (6.26)$$

and analogous insertions into the states (6.24). The insertion of the other linear combination of $H_B = 1$ fields $(A_1 \tilde{A}_2 - A_2 \tilde{A}_1)$ results in splitting the operator $\tilde{Q} \cdots Q$ into $\tilde{Q} \cdots Q \tilde{Q} \cdots Q$ by using the F-term equations. The latter operator is a “double-trace” operator which is subleading in the $1/N$ -expansion. Since the operator vanishes (modulo subleading terms) by using the F-term equations it is not a chiral primary operator (nor is it a “near chiral primary”) and is therefore ruled out.

Zero-momentum oscillators in the w_3 , $\bar{w}_3$ directions of $\tilde{\mathbb{R}}^4$ are absent from the open string spectrum [8], so the corresponding operators on the gauge theory side should be absent as well. This follows from the fact that the insertion of a ϕ (which has $H_B = 1$) can always be commuted to one of the endpoints by using the F-term equations, where it vanishes using the $0 = \tilde{Q}_I \phi$ F-term equation. Hence, such an operator is not chiral primary, and is therefore not protected. Similarly, insertions of $\phi^\dagger$ should not be allowed, but this is more difficult to show.

The number of bosonic zero modes obtained above is in agreement with the string theory result [8], where it was shown that on the string theory side there are six bosonic and four fermionic zero modes.

7 The $\mathcal{N} = 2$ $\text{SU} \times \text{SU} \times \cdots \times \text{SU} \times \text{SU}$ orientifold

The $\text{SU}(N) + 2\begin{smallmatrix} \square \\ \square \end{smallmatrix} \oplus 4\square$ theory studied in section 6 is the first in an infinite series of similar theories [12]. The orientifold group for these theories has the form $G = \mathbb{Z}_{2N_2}^{\text{orb}} \times \mathbb{Z}_2^{\text{ori}}$, where the pure orientifold part has the universal form $\mathbb{Z}_2^{\text{ori}} = \{1, \Omega(-1)^{F_L} R_{45}\}$, and the orbifold part is $\mathbb{Z}_{2N_2}^{\text{orb}} = \{1, \theta, \dots, \theta^{2N_2-1}\}$, where θ acts on the 6789 directions as $(z_1, z_2) \rightarrow (e^{\pi i/N_2} z_1, e^{-\pi i/N_2} z_2)$. For a particular choice of the orientifold projections acting on the D3-branes, the gauge group becomes [12]

$$\text{SU}(v_1) \times \cdots \times \text{SU}(v_{N_2}), \quad (7.1)$$

with the matter content (in $\mathcal{N} = 2$ language)

$$\begin{smallmatrix} \square \\ \square \end{smallmatrix}_1 + \bigoplus_{j=2}^{N_2} (\begin{smallmatrix} \square \\ \square \end{smallmatrix}_{j-1}, \overline{\begin{smallmatrix} \square \\ \square \end{smallmatrix}}_j) + \begin{smallmatrix} \square \\ \square \end{smallmatrix}_{N_2} + \bigoplus_{j=1}^{N_2} w_j \begin{smallmatrix} \square \\ \square \end{smallmatrix}_j. \quad (7.2)$$

The w_j 's are non-negative integers constrained by $\sum_{j=1}^{N_2} w_j = 4$. The v_j 's obey constraints arising from the vanishing of the beta-function(s) of the field theory [12]. The adS/CFT correspondence involves the large N_1 limit, where $N_1 \equiv v_1$ denotes the rank of the first factor of the gauge group (denoted by N in the previous section). It can be shown [27] that, to leading order in N_2/N_1 , the v_j 's corresponding to the different gauge group factors are equal: $v_j = N_1 [1 + \mathcal{O}(N_2/N_1)]$. Thus for $N_2 \ll N_1$ the gauge group is essentially $\text{SU}(N_1)^{N_2}$. In the following we will not assume that $N_2 \ll N_1$, although there are some motivations for making this assumption as will be discussed later in this section. The quantum numbers of the scalar fields of the $\text{SU} \times \cdots \times \text{SU}$ gauge theory are given in the table below.

CFT field	Representation	J_R	J_R^3	J_L^3	Δ	H_B
ϕ_r	adjoint_r	1	0	0	1	1
$\phi_r^\dagger$	adjoint_r	-1	0	0	1	1
$(A_1, (\tilde{A}^1)^\dagger)$	$\begin{smallmatrix} \square \\ \square \end{smallmatrix}_1$	0	$(\frac{1}{2}, -\frac{1}{2})$	$\frac{1}{2}$	1	(0,1)
$(\tilde{A}^1, (A_1)^\dagger)$	$\begin{smallmatrix} \square \\ \square \end{smallmatrix}_1$	0	$(\frac{1}{2}, -\frac{1}{2})$	$-\frac{1}{2}$	1	(1,2)
$(A_2, (\tilde{A}^2)^\dagger)$	$\begin{smallmatrix} \square \\ \square \end{smallmatrix}_{N_2}$	0	$(\frac{1}{2}, -\frac{1}{2})$	$-\frac{1}{2}$	1	(1,2)
$(\tilde{A}^2, (A_2)^\dagger)$	$\begin{smallmatrix} \square \\ \square \end{smallmatrix}_{N_2}$	0	$(\frac{1}{2}, -\frac{1}{2})$	$\frac{1}{2}$	1	(0,1)
$(B_p, C_p^\dagger)$	$(\overline{\begin{smallmatrix} \square \\ \square \end{smallmatrix}}_{p-1}, \begin{smallmatrix} \square \\ \square \end{smallmatrix}_p)$	0	$(\frac{1}{2}, -\frac{1}{2})$	$\frac{1}{2}$	1	(0,1)
$(C_p, B_p^\dagger)$	$(\begin{smallmatrix} \square \\ \square \end{smallmatrix}_{p-1}, \overline{\begin{smallmatrix} \square \\ \square \end{smallmatrix}}_p)$	0	$(\frac{1}{2}, -\frac{1}{2})$	$-\frac{1}{2}$	1	(1,2)
$(Q_r, \tilde{Q}_r^\dagger)$	$\frac{1}{2} w_r \begin{smallmatrix} \square \\ \square \end{smallmatrix}_r$	0	$(\frac{1}{2}, -\frac{1}{2})$	0	1	$(\frac{1}{2}, \frac{3}{2})$
$(\tilde{Q}_r, Q_r^\dagger)$	$\frac{1}{2} w_r \overline{\begin{smallmatrix} \square \\ \square \end{smallmatrix}}_r$	0	$(\frac{1}{2}, -\frac{1}{2})$	0	1	$(\frac{1}{2}, \frac{3}{2})$

Table 4: $\text{SU}(2)_R$ multiplets of scalar fields in the $\mathcal{N} = 2$ $\text{SU} \times \cdots \times \text{SU}$ orientifold theory.

In the table, r takes the values $r = 1, \dots, N_2$, whereas p ranges from 2 to N_2 . The quantum number J_L^3 in the table above denotes a $\text{U}(1)_L$ quantum number. Note that there is generically no $\text{SU}(2)_L$ symmetry (only when $N_2 = 1$ is there such a symmetry). The correspondence between our notation and that of ref. [13] is as follows: $J_R^3 \rightarrow J'$, $J_L^3 \rightarrow N_2 J$.

We will now show that there is a limit of the $SU \times \cdots \times SU$ theory discussed above in which the null direction x^- becomes compact, analogous to the limit discovered in refs. [13, 14]. Although the gauge group of the model is very similar to that of the unorientifolded model considered in refs. [13, 14], the matter content differs; in particular, there are no bifundamental fields connecting the last and first factors of the gauge group.

We consider only Penrose limit B, since it is in this limit that the compact null direction appears. Using the results of section 3, we find that the orbifold identifies (φ_1, w_2) with $(\varphi_1 + (\pi/N_2), e^{-i\pi/N_2} w_2)$. This identification along the geodesic translates into $x^+ \sim x^+ + (\pi/2N_2)$ and $x^- \sim x^- + (\pi R^2/2N_2)$. As observed in [13], if one take a scaling limit in which $2\pi R_- \equiv \pi R^2/2N_2 \sim (g_s N_1/N_2)^{1/2} \alpha'$ stays finite, the x^- direction becomes compact with radius R_- . Since we obtain a background with a compact null direction the string theory can be quantized using the discrete-light-cone-quantization (DLCQ) scheme, familiar from matrix theory [28].

Closed string sector

Since there is no representation connecting the last and first factor of the product gauge group, a gauge invariant operator cannot be constructed from a “loop” of $H_B = 0$ bifundamental fields B_p , as in ref. [13]. However, we may introduce the two $H_B = 0$ operators

$$\begin{aligned} \mathcal{A}_{a\bar{b}} &= (A_1 B_2 \cdots B_{N_2})_{a\bar{b}} \\ \tilde{\mathcal{A}}^{ab} &= (B_2 \cdots B_{N_2} \tilde{A}^2)^{ab} \end{aligned} \quad \begin{cases} a = 1, \dots, v_1 \\ \bar{b} = 1, \dots, v_{N_2} \end{cases} \quad (7.3)$$

From these, a gauge invariant operator

$$\text{tr}[(\mathcal{A}\tilde{\mathcal{A}}^T)^k] \quad (7.4)$$

can be constructed which may be identified with vacuum of the closed string sector. By the reasoning in ref. [13], k is the momentum labeling the different DLCQ sectors.

In ref. [13], the ground state corresponded to a “loop” of bifundamental fields, wrapping k times around the quiver/moose diagram. It was suggested that this was related to the T-dual type IIA picture of the gauge theory. In a similar way, the two operators in (7.3) can be identified with two half-circles of a quiver/moose diagram which are identified by the orientifold. This is reminiscent of the type IIA picture of the orientifold model, which contains two $O6^-$ planes located at diametrically opposite points of the elliptic circle, which reflect the circle onto itself [12]. It should be possible to connect these two pictures by applying a T-duality as in ref. [13]. (T-dualities of the pp-wave background have also been discussed in ref. [29]).

The operators corresponding to the string theory states may also be obtained using a cover space approach, as in ref. [13] and in sec. 6. Beginning with matrices Φ_i in the adjoint representation of an $SU(2N_v)$ theory, where $N_v = \sum_r v_r$, and projecting to the $SU(v_1) \times \cdots \times SU(v_{N_2})$ theory using a generalization [12] of eqs. (6.4), we obtain

$$\Phi_1 = \begin{pmatrix} \mathbf{B}^T & \mathbf{A}_1 \\ -\tilde{\mathbf{A}}^2 & \mathbf{B} \end{pmatrix}, \quad \Phi_2 = \begin{pmatrix} \mathbf{C} & \mathbf{A}_2 \\ \tilde{\mathbf{A}}^1 & \mathbf{C}^T \end{pmatrix}, \quad \Phi_3 = \begin{pmatrix} \phi & 0 \\ 0 & -\phi^T \end{pmatrix}, \quad (7.5)$$

where $\phi = \text{diag}(\phi_1, \phi_2, \dots, \phi_{N_2})$ and

$$\begin{aligned} \mathbf{A}_1 &= \text{diag}(A_1, 0, \dots, 0), & \tilde{\mathbf{A}}^1 &= \text{diag}(\tilde{A}^1, 0, \dots, 0), \\ \mathbf{A}_2 &= \text{diag}(0, \dots, 0, A_2), & \tilde{\mathbf{A}}^2 &= \text{diag}(0, \dots, 0, \tilde{A}^2), \end{aligned}$$

$$\mathbf{B} = \begin{pmatrix} 0 & B_2 & \cdots & 0 \\ 0 & 0 & \ddots & \vdots \\ \vdots & & \ddots & B_{N_2} \\ 0 & 0 & \cdots & 0 \end{pmatrix}, \quad \mathbf{C} = \begin{pmatrix} 0 & 0 & \cdots & 0 \\ C_2 & 0 & & 0 \\ \vdots & \ddots & \ddots & \vdots \\ 0 & \cdots & C_{N_2} & 0 \end{pmatrix}. \quad (7.6)$$

In terms of the Φ_i 's, the ground state in the closed string sector with lightcone momentum k can be written as

$$\text{tr}(\Phi_1^{2N_2k}), \quad (7.7)$$

which precisely coincides with eq. (7.4). Moreover, operators corresponding to string theory states with zero-momentum oscillators acting on the vacuum can be obtained via insertions of the $H_B = 1$ fields $(D_\mu \Phi_1)\Phi_1$, $\Phi_2\Phi_1$, $\Phi_2^\dagger\Phi_1$, $\Phi_3\Phi_1$, and $\Phi_3^\dagger\Phi_1$ into the vacuum operator, eq. (7.7).

Since we have a compact direction, winding modes are present in the string theory. These and the non-zero-momentum states can be constructed as in ref. [13]. As an example, a state with a single non-zero-momentum oscillator $a_n^\dagger$ in the w_3 direction acting on the vacuum can be obtained through an insertion of $\Phi_3 V^n$ into (7.7). (Due to the $\mathbb{Z}_2^{\text{ori}}$ projection, this vanishes if $n = 0$.) Here V is the diagonal matrix of phases $\sqrt{\omega} \text{diag}(\mathbb{1}_{v_1}, \omega \mathbb{1}_{v_2}, \dots, \omega^{N_2-1} \mathbb{1}_{v_{N_2}}; \omega^{2N_2-1} \mathbb{1}_{v_1}, \omega^{2N_2-2} \mathbb{1}_{v_2}, \dots, \omega^{N_2} \mathbb{1}_{v_{N_2}})$ with $\omega = \exp(\pi i/N_2)$ [12]. As explained in [13], such a state has winding number n . More generally, if the sum of the oscillator mode numbers does not vanish, then the state has non-zero winding number.

Open string sector

Since the geodesic in Penrose limit B lies in the orientifold (O7) fixed-plane $w_3 = 0$, which is coincident with the D7-branes, there is an open-string sector, in contrast to the model studied in ref. [13]. The model (7.1), (7.2) actually comprises several different models, due to the fact that there are several possible choices for the (non-negative) w_r 's satisfying $\sum_r w_r = 4$. The different models are distinguished by the fact that the fundamentals belong to different factors of the gauge group. To be able to treat all the different cases on the same footing we will use the following notation. Assuming that Q^I (or $\tilde{Q}_I$) for a particular I belongs to the r th factor of the gauge group, we can define

$$\begin{aligned} \mathcal{Q}_{\bar{a}}^I &= (Q^I B_{r+1} \cdots B_{N_2})_{\bar{a}}, \\ \tilde{\mathcal{Q}}_I^a &= (B_2 \cdots B_r \tilde{Q}_I)^a. \end{aligned} \quad (7.8)$$

From these, one can construct operators corresponding to the open string vacua, in analogy with (6.23), (6.24)

$$\mathcal{Q}^I \tilde{\mathcal{A}}^T (\mathcal{A} \tilde{\mathcal{A}}^T)^{(k-1)} A_1 \tilde{\mathcal{Q}}_K, \quad \mathcal{Q}^I \tilde{\mathcal{A}}^2 (\mathcal{A}^T \tilde{\mathcal{A}})^k (\mathcal{Q}^K)^T, \quad (\tilde{\mathcal{Q}}_I)^T (\mathcal{A}^T \tilde{\mathcal{A}})^k A_1 \tilde{\mathcal{Q}}_K. \quad (7.9)$$

As above, k is the non-negative integer lightcone momentum quantum number labeling the different DLCQ vacua. There are two kinds of open strings, as in section 6. (The operators (7.9) reduce to those in the previous section when $N_2 = 1$.)

Excited states in the open string sector are constructed by inserting $H_B = 1$ operators, as in section 6. Since there is no level matching condition in the open string sector, a single non-zero-momentum oscillator acting on the vacuum state is allowed.

Finally, we will discuss the implications of making the additional assumption $N_2 \ll N_1$. For the original model [2] the 't Hooft coupling is $\lambda' = g_{\text{YM}}^2 N / J^2$. It was recently argued [23] that the effective genus counting parameter in the Yang-Mills theory is $g_2^2 = J^4 / N^2$, and the effective coupling between a wide class of excited states is $g_2 \sqrt{\lambda'}$ (see ref. [23] for further details; see also [30] for some refinements). Translating these expressions to our case one finds that both the expansion parameters are small provided $N_2 \ll N_1$, thus from this perspective it is natural to make the additional assumption $N_2 \ll N_1$. Note that this assumption corresponds to a large radius approximation for the compact circle.

8 $\mathcal{N} = 2$ $\text{Sp}(2N) \times \text{Sp}(2N)$ **with** $(\square, \square) \oplus 2(1, \square) \oplus 2(\square, 1)$

The $\mathcal{N} = 2$ $\text{Sp}(2N) \times \text{Sp}(2N)$ theory with an $\mathcal{N} = 2$ hypermultiplet in the bifundamental representation $(\square, \square)$ and two $\mathcal{N} = 2$ hypermultiplets in the fundamental representation of each factor of the gauge group contains (in $\mathcal{N} = 1$ language) two chiral multiplets $\phi_{1a}{}^b$ and $\phi_{2\bar{a}}{}^{\bar{b}}$ in the adjoint representation, one for each $\text{Sp}(2N)$ factor. In matrix notation, these fields satisfy

$$\phi_1 = J_1 \phi_1^T J_1, \quad \phi_2 = J_2 \phi_2^T J_2, \quad (8.1)$$

where $J^{ab} = J_1$ and $J_{ab} = J_1^{-1} = -J_1$ is the symplectic unit of the first $\text{Sp}(2N)$ factor, used to raise and lower indices (and similarly for $J^{\bar{a}\bar{b}} = J_2$). The theory also contains two chiral multiplets in the $(\square, \square)$ representation; for convenience, in writing operators in matrix notation, we represent these with two chiral multiplets $A_{ia}{}^{\bar{b}}$ ($i = 1, 2$) in the $(\square, \bar{\square})$ representation of $\text{Sp}(2N) \times \text{Sp}(2N)$ and two chiral multiplets $B_{i\bar{a}}{}^b$ ($i = 1, 2$) in the $(\bar{\square}, \square)$ representation, with the constraint

$$B_i = -J_2 A_i^T J_1. \quad (8.2)$$

In this paper, both (A_1, A_2) and (B_1, B_2) are in the $\square$ of $\text{SU}(2)_L$. (The indices may be raised with the ϵ tensor, so that $(B^1, B^2) \equiv (B_2, -B_1)$ is in the $\bar{\square}$ of $\text{SU}(2)_L$. In ref. [31] and many other papers, (B^1, B^2) is written as (B_1, B_2) .)

Finally, the theory also contains four chiral multiplets in the $(\square, 1)$ and $(1, \square)$ representations, denoted by Q_{1a}^I and $Q_{2\bar{a}}^I$ ($I = 1, \dots, 4$), respectively. For convenience, we also introduce fields $\tilde{Q}_{1I}^a$ and $\tilde{Q}_{2I}^{\bar{a}}$ in the $(\bar{\square}, 1)$ and $(1, \bar{\square})$ representations, related to the above fields by [31]

$$\tilde{Q}_{1I}^T = -g_{IJ} J_1 Q_1^J, \quad \tilde{Q}_{2I}^T = -g_{IJ} J_2 Q_2^J, \quad (8.3)$$

where g_{IJ} is the metric for either of the two factors of the $\text{SO}(4) \times \text{SO}(4)$ flavor symmetry group. The quantum numbers of the scalar fields are displayed in the table below.

CFT field	$\text{Sp}(2N) \times \text{Sp}(2N)$	J_R	J_R^3	J_L^3	Δ	H_A	H_B
ϕ_1	(adjoint, 1)	1	0	0	1	0	1
$\phi_1^\dagger$	(adjoint, 1)	-1	0	0	1	2	1
ϕ_2	(1, adjoint)	1	0	0	1	0	1
$\phi_2^\dagger$	(1, adjoint)	-1	0	0	1	2	1
$(A_1, B_1^\dagger)$	$(\square, \overline{\square})$	0	$(\frac{1}{2}, -\frac{1}{2})$	$\frac{1}{2}$	1	(1, 1)	(0, 1)
$(A_2, B_2^\dagger)$	$(\square, \overline{\square})$	0	$(\frac{1}{2}, -\frac{1}{2})$	$-\frac{1}{2}$	1	(1, 1)	(1, 2)
$(B_1, A_1^\dagger)$	$(\overline{\square}, \square)$	0	$(\frac{1}{2}, -\frac{1}{2})$	$\frac{1}{2}$	1	(1, 1)	(0, 1)
$(B_2, A_2^\dagger)$	$(\overline{\square}, \square)$	0	$(\frac{1}{2}, -\frac{1}{2})$	$-\frac{1}{2}$	1	(1, 1)	(1, 2)
$(Q_1, \tilde{Q}_1^\dagger)$	$(\square, 1)$	0	$(\frac{1}{2}, -\frac{1}{2})$	0	1	(1, 1)	$(\frac{1}{2}, \frac{3}{2})$
$(\tilde{Q}_1, Q_1^\dagger)$	$(\overline{\square}, 1)$	0	$(\frac{1}{2}, -\frac{1}{2})$	0	1	(1, 1)	$(\frac{1}{2}, \frac{3}{2})$
$(Q_2, \tilde{Q}_2^\dagger)$	$(1, \square)$	0	$(\frac{1}{2}, -\frac{1}{2})$	0	1	(1, 1)	$(\frac{1}{2}, \frac{3}{2})$
$(\tilde{Q}_2, Q_2^\dagger)$	$(1, \overline{\square})$	0	$(\frac{1}{2}, -\frac{1}{2})$	0	1	(1, 1)	$(\frac{1}{2}, \frac{3}{2})$

Table 5: $\text{SU}(2)_R$ multiplets of scalar fields in $\mathcal{N} = 2$ $\text{Sp}(2N) \times \text{Sp}(2N)$ with $(\square, \square) \oplus 2(1, \square) \oplus 2(\square, 1)$.

The superpotential for this theory is

$$\mathcal{W}_{\mathcal{N}=2} = \text{tr} \left[\phi_1 (A_1 B^1 + A_2 B^2) - \phi_2 (B^1 A_1 + B^2 A_2) \right] + \tilde{Q}_{1I} \phi_1 Q_1^I + \tilde{Q}_{2I} \phi_2 Q_2^I, \quad (8.4)$$

giving rise to the F-term equations

$$\begin{aligned} A_1 B_2 - A_2 B_1 + Q_1^I \tilde{Q}_{1I} &= 0, & A_i \phi_2 - \phi_1 A_i &= 0, & \phi_1 Q_1^I &= 0, \\ B_1 A_2 - B_2 A_1 + Q_2^I \tilde{Q}_{2I} &= 0, & B_i \phi_1 - \phi_2 B_i &= 0, & \phi_2 Q_2^I &= 0. \end{aligned} \quad (8.5)$$

The $\mathcal{N} = 2$ $\text{Sp}(2N) \times \text{Sp}(2N)$ gauge theory just discussed arises as the theory on a stack of D3-branes in the same IIB background that gave rise to the $\text{SU}(N)$ with $2\overline{\square} \oplus 4\square$ theory discussed in section 6, namely, flat space modded out by

$$G = \mathbb{Z}_2^{\text{orb}} \times \mathbb{Z}_2^{\text{ori}}. \quad (8.6)$$

As in that case, the fixed point set of $\mathbb{Z}_2^{\text{ori}}$ is the hyperplane $z_3 = 0$, which corresponds to the position of the O7-plane and the D7-branes, while the fixed point set of $\mathbb{Z}_2^{\text{orb}}$ is the six-dimensional hyperplane $z_1 = z_2 = 0$.

The fields of the $\mathcal{N} = 2$ $\text{Sp}(2N) \times \text{Sp}(2N)$ theory can be obtained via a projection (different from the one in section 6) from the $\mathcal{N} = 4$ $\text{SU}(4N)$ gauge theory. The independent generators of the orientifold group are realized as the $4N \times 4N$ matrices [12]

$$\gamma_\theta = \begin{pmatrix} \mathbb{1}_{2N} & 0 \\ 0 & -\mathbb{1}_{2N} \end{pmatrix}, \quad \gamma_{\Omega'} = \begin{pmatrix} J & 0 \\ 0 & J \end{pmatrix}. \quad (8.7)$$

The orbifold projection

$$\Phi_{1,2} = -\gamma_\theta \Phi_{1,2} \gamma_\theta^{-1}, \quad \Phi_3 = \gamma_\theta \Phi_3 \gamma_\theta^{-1}, \quad (8.8)$$

restricts the adjoint fields, Φ_i , of the $\mathcal{N} = 4$ theory to be of the form

$$\Phi_1 = \begin{pmatrix} 0 & A_1 \\ -B^2 & 0 \end{pmatrix} = \begin{pmatrix} 0 & A_1 \\ B_1 & 0 \end{pmatrix}, \quad \Phi_2 = \begin{pmatrix} 0 & A_2 \\ B^1 & 0 \end{pmatrix} = \begin{pmatrix} 0 & A_2 \\ B_2 & 0 \end{pmatrix}, \quad \Phi_3 = \begin{pmatrix} \phi_1 & 0 \\ 0 & \phi_2 \end{pmatrix}. \quad (8.9)$$

The orientifold projection

$$\Phi_{1,2} = \gamma_{\Omega'} \Phi_{1,2}^T \gamma_{\Omega'}^{-1}, \quad \Phi_3 = -\gamma_{\Omega'} \Phi_3^T \gamma_{\Omega'}^{-1}, \quad (8.10)$$

is equivalent to the constraints (8.1) and (8.2).

The action of G on the coordinates of $adS_5 \times S^5$ is as follows. $\mathbb{Z}_2^{\text{orb}}$ acts as $\varphi_1 \rightarrow \varphi_1 + \pi$, $\varphi_2 \rightarrow \varphi_2 + \pi$ and $\mathbb{Z}_2^{\text{ori}}$ acts as $\varphi_3 \rightarrow \varphi_3 + \pi$. String theory on this background is dual to the above $\text{Sp} \times \text{Sp}$ gauge theory [10, 11]. Below we discuss the pp-wave limits of this correspondence.

Penrose limit A

In the first Penrose limit, the generator of $\mathbb{Z}_2^{\text{ori}}$ simply produces a translation along the geodesic: $\varphi_3 \rightarrow \varphi_3 + \pi$ (together with the action of $\Omega(-1)^{F_L}$) with no action on the transverse coordinates w_1 and w_2 (3.9). There is no orientifold fixed plane and therefore no open string sector in this Penrose limit, but $\mathbb{Z}_2^{\text{ori}}$ projects out half of the discrete values of J_A that would be allowed in the unprojected case. The generator of $\mathbb{Z}_2^{\text{orb}}$ acts solely on the transverse coordinates as $w_1 \rightarrow -w_1$, $w_2 \rightarrow -w_2$. The geodesic lies in the orbifold fixed plane, resulting in a twisted sector of the string theory in this pp-wave background.

Untwisted and twisted sectors

As in the case of the $\text{SU}(N) + 2\Box \oplus 4\Box$ theory in section 6, the operators in the untwisted and twisted sectors are easily obtained through projections of the cover space fields.

The ground state in the untwisted sector corresponds to the operator $\text{tr}[\Phi_3^{J_A}]$ whereas the twisted sector ground state corresponds to the operator $\text{tr}[\gamma_\theta \Phi_3^{J_A}]$. The $\mathbb{Z}_2^{\text{ori}}$ projection (8.10) implies $\text{tr}[\gamma_\theta^q \Phi_3^{J_A}] = (-)^{J_A} \text{tr}[\gamma_\theta^q (\Phi_3^T)^{J_A}] = (-)^{J_A} \text{tr}[\gamma_\theta^q \Phi_3^{J_A}]$, which vanishes unless $J_A = 2n$ for both the untwisted and twisted sectors. Thus, the $H_A = 0$ operators

$$\text{tr} [\phi_1^{2n} \pm \phi_2^{2n}], \quad (8.11)$$

correspond to the vacuum states in the untwisted (+) and twisted (−) sectors of the string theory.

The $H_A = 1$ gauge theory operators $\text{tr}[(D_\mu \Phi_3) \Phi_3^{2n+1}]$ and $\text{tr}[\gamma_\theta (D_\mu \Phi_3) \Phi_3^{2n+1}]$ or, explicitly,

$$\text{tr} [(D_\mu \phi_1) \phi_1^{2n+1} \pm (D_\mu \phi_2) \phi_2^{2n+1}], \quad (8.12)$$

correspond to string states in which a single zero-momentum oscillator $a_0^{\dagger\mu}$ in one of the directions of $\mathbb{R}^4$ acts on the light-cone vacuum in the untwisted (+) and twisted (−) sectors respectively.

Since the zero-momentum oscillators $a_0^{\dagger I}$ in the transverse $\tilde{\mathbb{R}}^4$ directions (3.9) are odd under $\mathbb{Z}_2^{\text{orb}}$, only pairs of such oscillators can act on the light-cone vacuum. The $H_A = 2$ gauge theory operators corresponding to the states involving two oscillators in the untwisted

sector involve the insertion of a pair of $H_A = 1$ fields $\Upsilon_I = (\Phi_1, \Phi_1^\dagger, \Phi_2, \Phi_2^\dagger)$ into the ground state operator (the insertion of a single $J_A = 1$ operator into the ground state operator gives zero using (8.9)):

$$\sum_{m=0}^{J_A} \text{tr} \left[\Upsilon_I \Phi_3^m \Upsilon_K \Phi_3^{J_A-m} \right], \quad (8.13)$$

where we average over the relative position of the Υ 's [2, 22, 23]. Eq. (8.13) vanishes unless J_A is even. Using (8.9), together with

$$\Phi_1^\dagger = \begin{pmatrix} 0 & B_1^{\dagger 1} \\ A_1^{\dagger 1} & 0 \end{pmatrix} = \begin{pmatrix} 0 & B_2^\dagger \\ A_2^\dagger & 0 \end{pmatrix}, \quad \Phi_2^\dagger = \begin{pmatrix} 0 & B_1^{\dagger 2} \\ A_1^{\dagger 2} & 0 \end{pmatrix} = \begin{pmatrix} 0 & -B_1^\dagger \\ -A_1^\dagger & 0 \end{pmatrix}, \quad (8.14)$$

we obtain the set of ten $H_A = 2$ operators

$$\begin{aligned} \sum_m \text{tr} \left[A_{(i} \phi_2^m B_{j)} \phi_1^{2n-m} \right], \quad \sum_m \text{tr} \left[B_{(i}^\dagger \phi_2^m A_{j)}^\dagger \phi_1^{2n-m} \right], \\ \sum_m \text{tr} \left[A_i \phi_2^m A_j^\dagger \phi_1^{2n-m} + B_j^\dagger \phi_2^m B_i \phi_1^{2n-m} \right], \end{aligned} \quad (8.15)$$

in agreement with string theory expectations, cf. (6.12).

The oscillators in the directions w_1 and w_2 are half-integer moded in the twisted sector [24, 25, 20, 26] so there exist no zero-momentum oscillators in this sector. The operators that would have corresponded to such oscillators acting on the twisted-sector vacuum

$$\sum_m \text{tr} \left[\gamma_\theta \Upsilon_I \Phi_3^m \Upsilon_K \Phi_3^{J_A-m} \right], \quad (8.16)$$

must therefore not be chiral primary operators in the gauge theory. Consider for example

$$\sum_m \text{tr} \left[A_{[i} \phi_2^m B_{j]} \phi_1^{J_A-m} \right], \quad (8.17)$$

Using the F-term equation $A_i \phi_2 = \phi_1 A_i$, A_i can be shifted through $(\phi_2)^m$. Then $A_{[i} B_{j]}$ can be replaced by Q_1 's, using $A_1 B_2 - A_2 B_1 + Q_1^I Q_{1I} = 0$. Finally, using $\phi_1 Q_1^I = 0$, the operator vanishes and is therefore not a chiral primary. The other operators in (8.16) are similarly expected to be ruled out.

Operators corresponding to non-zero-momentum oscillators can be constructed in analogy with the constructions in refs. [25, 20, 26].

Penrose limit B

In limit B, the geometric action of $\mathbb{Z}_2^{\text{ori}}$, $\varphi_3 \rightarrow \varphi_3 + \pi$, produces a reflection of the transverse coordinate $w_3 \rightarrow -w_3$. The geodesic lies in the orientifold (O7) plane $w_3 = 0$, and the coincident D7-branes give rise to an open-string sector. The generator of $\mathbb{Z}_2^{\text{orb}}$ produces a shift $\varphi_1 \rightarrow \varphi_1 + \pi$ halfway around the geodesic, accompanied by a reflection of the coordinates transverse to the O7-plane: $w_2 \rightarrow -w_2$. Since the $\mathbb{Z}_2^{\text{orb}}$ action has no fixed points, there is no twisted sector in this Penrose limit, however $\mathbb{Z}_2^{\text{orb}}$ projects out half of the discrete values of J_B that would be allowed in the unprojected case.

Closed string sector

The vacuum state of the closed string sector corresponds to the $H_B = 0$ operator $\text{tr}[\Phi_1^{J_B}]$. Due to the $\mathbb{Z}_2^{\text{orb}}$ projection on Φ_1 , this is only nonvanishing when $J_B = 2n$,

$$\text{tr}[\Phi_1^{2n}] = \text{tr}[(A_1 B_1)^n]. \quad (8.18)$$

String states in which a zero-momentum oscillator $a_0^{\dagger\mu}$ in one of the directions of $\mathbb{R}^4$ acts on the light-cone vacuum correspond to the $H_B = 1$ gauge theory operators

$$\text{tr}[(D_\mu \Phi_1) \Phi_1^{2n+1}] = \text{tr}[D_\mu (A_1 B_1) (A_1 B_1)^n]. \quad (8.19)$$

String states in which a zero-momentum oscillator $a_0^\dagger, \bar{a}_0^\dagger$ in the $w_2, \bar{w}_2$ directions of $\tilde{\mathbb{R}}^4$ acts on the light-cone vacuum corresponds to insertions of the $H_B = 1$ fields $\Phi_2 \Phi_1, \Phi_2^\dagger \Phi_1$ into the vacuum state operator:

$$\begin{aligned} \text{tr}[\Phi_2 \Phi_1^{2n+1}] &= \text{tr}[(A_2 B_1 + A_1 B_2) (A_1 B_1)^n], \\ \text{tr}[\Phi_2^\dagger \Phi_1^{2n+1}] &= \text{tr}[(B_1^\dagger B_1 + A_1 A_1^\dagger) (A_1 B_1)^n]. \end{aligned} \quad (8.20)$$

The linear combination $A_2 B_1 - A_1 B_2$ is ruled out since it is not a chiral primary; see section 6 for further details.

The zero-momentum oscillators in the w_3 ($\bar{w}_3$) directions of $\tilde{\mathbb{R}}^4$ are odd under $\mathbb{Z}_2^{\text{ori}}$, so there are no states corresponding to single oscillators acting on the vacuum. Correspondingly, the gauge theory operator $\text{tr}[\Phi_3 \Phi_1^{J_B}]$, resp. $\text{tr}[\Phi_3^\dagger \Phi_1^{J_B}]$, vanishes, as can be seen by taking the transpose of $[\phi_1 (A_1 B_1)^n]$, resp. $[\phi_1^\dagger (A_1 B_1)^n]$, and using (8.1) and (8.2).

Open string sector

Since the geodesic lies in the orientifold (O7) fixed-plane $w_3 = 0$ which is also the location of the D7-branes, there is an open-string sector. The gauge symmetry of the D7-branes in the presence of the O7-plane, $\text{SO}(8)_F$, is broken to $\text{SO}(4) \times \text{SO}(4)$ by the $\mathbb{Z}_2^{\text{orb}}$ projection. The ground state of the open string sector, which transforms in the adjoint **28** of $\text{SO}(8)_F$, decomposes into

$$\mathbf{28} \rightarrow (\mathbf{6}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{6}) \oplus (\mathbf{4}, \mathbf{4}). \quad (8.21)$$

The $\mathbb{Z}_2^{\text{orb}}$ projection correlates the states in the $(\mathbf{6}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{6})$ representation with operators in odd-dimensional representations of $\text{SU}(2)_L$ [11]. These open string vacuum states therefore correspond to gauge theory operators with an even number of A 's and B 's. The natural candidates are

$$Q_1^I J_1 (A_1 B_1)^n Q_1^K, \quad Q_2^I J_2 (B_1 A_1)^n Q_2^K. \quad (8.22)$$

These operators have $H_B = 1$, reflecting the non-vanishing zero point energy for the open string sector [8]. The antisymmetry of the flavor indices I, K is enforced by the antisymmetry of the matrices $J_1 (A_1 B_1)^n$ and $J_2 (B_1 A_1)^n$ (which follows from eq. (8.2)) and guarantees that the operators (8.22) belong to the correct representation of the flavor group.

The $\mathbb{Z}_2^{\text{orb}}$ projection correlates the states in the $(\mathbf{4}, \mathbf{4})$ representation with operators in even-dimensional representations of $\text{SU}(2)_L$ [11]. These open string vacuum states therefore

correspond to the gauge theory operators with an odd numbers of A 's and B 's. The natural candidate is

$$Q_1^I J_1 (A_1 B_1)^n A_1 Q_2^K, \quad (8.23)$$

which has $H_B = 1$ and belongs to the $(\mathbf{4}, \mathbf{4})$ representation of the flavor group. (The operator $Q_2^K J_2 B_1 (A_1 B_1)^n Q_1^I$ is equivalent to (8.23) using the orientifold projections (8.2).)

Zero-momentum oscillators in the $w_2, \bar{w}_2$ directions of $\tilde{\mathbb{R}}^4$ acting on the open string vacuum correspond to the insertion of the $H_B = 1$ fields $A_2 B_1 + A_1 B_2$ and $B_1^\dagger B_1 + A_1 A_1^\dagger$ into the operators (8.22) and (8.23).

Zero-momentum oscillators in the $w_3, \bar{w}_3$ directions of $\tilde{\mathbb{R}}^4$ (which are the directions transverse to the D7-branes) are absent from the open string spectrum [8], so the corresponding operators on the gauge theory side should be absent as well. This follows from the fact that, as in section 6, the insertion of $\phi_{1,2}$ (which has $H_B = 1$) can be shown to give rise to an operator that vanishes by the F-term equations. It is therefore not a chiral primary operator and is consequently ruled out, in agreement with the string theory result [8].

9 The $\mathcal{N} = 2$ $\text{Sp} \times \text{SU} \times \cdots \times \text{SU} \times \text{Sp}$ orientifold

The $\text{Sp}(2N) \times \text{Sp}(2N)$ theory discussed in section 8 is the first in an infinite series of conformal models with gauge groups $\text{Sp} \times \text{SU} \times \cdots \times \text{SU} \times \text{Sp}$ [12]. The orientifold group for these theories has the form $G = \mathbb{Z}_{2k}^{\text{orb}} \times \mathbb{Z}_2^{\text{ori}}$. As before, $\mathbb{Z}_2^{\text{ori}} = \{\mathbf{1}, \Omega(-1)^{F_L} R_{45}\}$ and $\mathbb{Z}_{2N_2}^{\text{orb}} = \{1, \theta, \dots, \theta^{2N_2-1}\}$. The action of the orbifold on the coordinates transverse to the D3-branes is the same as in section 7. For a particular choice of the orientifold projection acting on the D3-branes (different from the one in section 7) the gauge group becomes [12]

$$\text{Sp}(v_0) \times \text{SU}(v_1) \times \cdots \times \text{SU}(v_{N_2-1}) \times \text{Sp}(v_{N_2}), \quad (9.1)$$

and the matter content is (in $\mathcal{N} = 2$ language)

$$\bigoplus_{j=1}^{N_2} (\square_{j-1}, \bar{\square}_j) + \frac{1}{2} w_0 \square_0 + \bigoplus_{j=1}^{N_2-1} w_j \square_j + \frac{1}{2} w_{N_2} \square_{N_2}. \quad (9.2)$$

The w_j 's are non-negative integers constrained by the equation $\frac{1}{2}(w_0 + w_{N_2}) + \sum_{j=1}^{N_2-1} w_j = 4$. The v_j 's obey constraints arising from the vanishing of the beta-function(s) of the field theory [12]. To leading order in N_2/N_1 , where $N_1 (\equiv v_0/2)$ equals the rank of the first factor of the product gauge group, the v_j 's corresponding to each of the group factors are equal and take the value $2N_1$ [27], and the gauge group is essentially $\text{Sp}(2N_1) \times \text{SU}(2N_1)^{N_2-1} \times \text{Sp}(2N_1)$. We will not make any assumptions about the magnitude of the ratio N_2/N_1 ; see, however, the discussion in section 7. The quantum numbers of the scalar fields of the $\text{Sp} \times \text{SU} \times \cdots \times \text{SU} \times \text{Sp}$ gauge theory (9.1) are given in the table below.

CFT field	Representation	J_R	J_R^3	J_L^3	Δ	H_B
ϕ_r	adjoint_r	1	0	0	1	1
$\phi_r^\dagger$	adjoint_r	-1	0	0	1	1
$(A_p, (B^p)^\dagger)$	$(\square_{p-1}, \overline{\square}_p)$	0	$(\frac{1}{2}, -\frac{1}{2})$	$\frac{1}{2}$	1	(0,1)
$(B^p, (A_p)^\dagger)$	$(\overline{\square}_{p-1}, \square_p)$	0	$(\frac{1}{2}, -\frac{1}{2})$	$-\frac{1}{2}$	1	(1,2)
$(Q_r, \tilde{Q}_r^\dagger)$	$\frac{1}{2}w_r \square_r$	0	$(\frac{1}{2}, -\frac{1}{2})$	0	1	$(\frac{1}{2}, \frac{3}{2})$
$(\tilde{Q}_r, Q_r^\dagger)$	$\frac{1}{2}w_r \overline{\square}_r$	0	$(\frac{1}{2}, -\frac{1}{2})$	0	1	$(\frac{1}{2}, \frac{3}{2})$

Table 6: $\text{SU}(2)_R$ multiplets of scalar fields in the $\mathcal{N} = 2 \text{ Sp} \times \text{SU} \times \cdots \times \text{SU} \times \text{Sp}$ theory.

In the table, r takes the values $r = 0, \dots, N_2$, whereas p ranges from 1 to N_2 . The quantum number J_L^3 in the table above denotes a $\text{U}(1)_L$ quantum number; generically there is no $\text{SU}(2)_L$ symmetry (except when $N_2 = 1$).

Scaling N_1 and N_2 to infinity together in Penrose limit B, as in sec. 7, we obtain a background with a compact null direction, in which the string theory can be quantized using the DLCQ scheme, as in ref. [13].

Closed string sector

Since there is no bifundamental representation connecting the last and first factors of the gauge group, one cannot construct a gauge invariant operator as a “loop” of $H_B = 0$ bifundamental fields. However, we may define

$$\mathcal{A}_a^{\bar{b}} = (A_1 A_2 \cdots A_{N_2})_a^{\bar{b}}, \quad a = 1, \dots, v_0, \quad \bar{b} = 1, \dots, v_{N_2}, \quad (9.3)$$

and then use the symplectic units J_1 and J_2 , of $\text{Sp}(v_0)$ and $\text{Sp}(v_{N_2})$ respectively, to define $\mathcal{B}_a^{\bar{b}}$ by $\mathcal{B} = J_2 \mathcal{A}^T J_1$. In terms of these two operators, a gauge invariant $H_B = 0$ operator may be constructed

$$\text{tr}[(\mathcal{A}\mathcal{B})^k], \quad (9.4)$$

which may be identified with the closed string ground state. Here k is an arbitrary non-negative integer which labels the DLCQ momentum states as in [13].

The pictorial interpretation of this operator is as a “loop” around a quiver/moose diagram, with two half-circles related via the orientifold projection (again reminiscent of the type IIA picture of the gauge theory).

As in section 7, one can write the gauge invariant operators corresponding to the vacuum states (9.4) as well as the excited states in terms of cover space fields.

Open string sector

The open string vacuum is constructed out of a string of $H_B = 0$ operators with two Q ’s at the ends, exactly as in section 7. Assuming that Q^I (or $\tilde{Q}^I$) for a particular I belongs to the r th factor of the gauge group, we can define

$$\begin{aligned} \mathcal{Q}_a^I &= (A_1 \cdots A_r Q^I)_a, \\ \tilde{\mathcal{Q}}_I^{\bar{a}} &= (\tilde{Q}^I A_{r+1} \cdots A_{N_2})^{\bar{a}}. \end{aligned} \quad (9.5)$$

From these, one can construct operators corresponding to the open string vacua, in analogy with (8.22), (8.23)

$$(\mathcal{Q}^I)^T J_1 (\mathcal{AB})^k \mathcal{Q}^K, \quad \tilde{\mathcal{Q}}_I (\mathcal{BA})^k J_2 (\tilde{\mathcal{Q}}_K)^T, \quad \tilde{\mathcal{Q}}_I \mathcal{B} (\mathcal{AB})^k \mathcal{Q}^K. \quad (9.6)$$

Further details can be found in section 7.

In addition to the series of models studied in this section and in section 7, there is another infinite series of models containing D7 branes, with gauge group $\mathrm{Sp} \times \mathrm{SU} \times \cdots \times \mathrm{SU} \times \mathrm{SU}$ [12]. These models can also be treated using the methods of this paper, and possess both a closed and open string sector in the Penrose limit.

In ref. [12] four additional infinite series of models were constructed that do not contain D7-branes. It should also be possible to analyze the pp-wave limits of these models.

10 $\mathcal{N} = 1$ $\mathrm{Sp}(2N) \times \mathrm{Sp}(2N)$ **with** $2(\square, \square) \oplus 4(1, \square) \oplus 4(\square, 1)$

In the previous sections, we have discussed various $\mathcal{N} = 2$ theories and their associated Penrose limits. It is of interest to extend these results to $\mathcal{N} = 1$ theories. The pp-wave limit associated with the duality between the $\mathcal{N} = 1$ $\mathrm{SU}(N) \times \mathrm{SU}(N)$ theory and string theory on $adS_5 \times T^{11}$ [32] was treated in refs. [24, 33, 34]. Orbifolds of this theory and their Penrose limits were discussed in ref. [35]. Various $\mathcal{N} = 1$ orbifolds of $adS_5 \times S^5$ have been discussed in ref. [33, 17, 35]; recently non-supersymmetric models have also been discussed [36].

In this section we discuss a particular orientifolded version of the $adS_5 \times T^{11}$ model and its Penrose limit. By studying this model we are able to test the ideas of ref. [8] in a more complicated example. The orientifold we study was previously discussed in ref. [27, 31, 37]. The field theory dual is an $\mathcal{N} = 1$ $\mathrm{Sp}(2N) \times \mathrm{Sp}(2N)$ field theory with matter content (in $\mathcal{N} = 1$ language) $2(\square, \square) \oplus 4(\square, 1) \oplus 4(1, \square)$. As in ref. [31] and in sec. 8, for notational clarity the bifundamental chiral multiplets are denoted using the doubled set of fields $A_{ia}^{\bar{b}}$ ($i = 1, 2$) and $B_{\bar{a}}^i{}^b$ ($i = 1, 2$), with the constraint⁴

$$B^1 = -J_2 A_2^T J_1, \quad B^2 = J_2 A_1^T J_1. \quad (10.1)$$

The chiral multiplets in the $(\square, 1)$ and $(1, \square)$ representations will be denoted by Q_{1a}^I and $Q_{2\bar{a}}^I$ ($I = 1, \dots, 4$), respectively. For convenience, we also introduce fields $\tilde{Q}_{1I}^a$ and $\tilde{Q}_{2I}^{\bar{a}}$, related to Q_{1a}^I and $Q_{2\bar{a}}^I$ by [31]

$$\tilde{Q}_{1I}^T = -g_{IJ} J_1 Q_1^J, \quad \tilde{Q}_{2I}^T = -g_{IJ} J_2 Q_2^J, \quad (10.2)$$

where g_{IJ} is the metric for either of the two factors of the $\mathrm{SO}(4) \times \mathrm{SO}(4)$ flavor symmetry group. The quantum numbers of the scalar fields and their complex conjugates are displayed in the table below.

⁴In ref. [31], (B^1, B^2) was written as (B_1, B_2) .

Field	Representation	J_R	J_R^3	Δ	H	Field	Representation	J_R	J_R^3	Δ	H
A_1	$(\square, \overline{\square})$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{3}{4}$	0	$(A_1)^\dagger$	$(\overline{\square}, \square)$	$-\frac{1}{2}$	$-\frac{1}{2}$	$\frac{3}{4}$	$\frac{3}{2}$
A_2	$(\square, \overline{\square})$	$\frac{1}{2}$	$-\frac{1}{2}$	$\frac{3}{4}$	1	$(A_2)^\dagger$	$(\overline{\square}, \square)$	$-\frac{1}{2}$	$\frac{1}{2}$	$\frac{3}{4}$	$\frac{1}{2}$
B^1	$(\overline{\square}, \square)$	$\frac{1}{2}$	$-\frac{1}{2}$	$\frac{3}{4}$	1	$(B^1)^\dagger$	$(\square, \overline{\square})$	$-\frac{1}{2}$	$\frac{1}{2}$	$\frac{3}{4}$	$\frac{1}{2}$
B^2	$(\overline{\square}, \square)$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{3}{4}$	0	$(B^2)^\dagger$	$(\square, \overline{\square})$	$-\frac{1}{2}$	$-\frac{1}{2}$	$\frac{3}{4}$	$\frac{3}{2}$
Q_1^I	$(\square, 1)$	$\frac{1}{2}$	0	$\frac{3}{4}$	$\frac{1}{2}$	$(Q_1^I)^\dagger$	$(\overline{\square}, 1)$	$-\frac{1}{2}$	0	$\frac{3}{4}$	$\frac{3}{2}$
$\tilde{Q}_{1I}$	$(\overline{\square}, 1)$	$\frac{1}{2}$	0	$\frac{3}{4}$	$\frac{1}{2}$	$(\tilde{Q}_{1I})^\dagger$	$(\square, 1)$	$-\frac{1}{2}$	0	$\frac{3}{4}$	$\frac{3}{2}$
Q_2^I	$(1, \square)$	$\frac{1}{2}$	0	$\frac{3}{4}$	$\frac{1}{2}$	$(Q_2^I)^\dagger$	$(1, \overline{\square})$	$-\frac{1}{2}$	0	$\frac{3}{4}$	$\frac{3}{2}$
$\tilde{Q}_{2I}$	$(1, \overline{\square})$	$\frac{1}{2}$	0	$\frac{3}{4}$	$\frac{1}{2}$	$(\tilde{Q}_{2I})^\dagger$	$(1, \square)$	$-\frac{1}{2}$	0	$\frac{3}{4}$	$\frac{3}{2}$

Table 7: Scalar fields in the $\mathcal{N} = 1$ $\text{Sp}(2N) \times \text{Sp}(2N)$ theory with $2(\square, \square) \oplus 4(1, \square) \oplus 4(\square, 1)$.

The superpotential for the $\mathcal{N} = 1$ $\text{Sp}(2N) \times \text{Sp}(2N)$ theory is given by [31]

$$\begin{aligned} \mathcal{W}_{\mathcal{N}=1} = & - \left[\text{tr}(A_1 B^1 A_2 B^2 - B^1 A_1 B^2 A_2) + \frac{1}{2} \tilde{Q}_{1I} Q_1^I \tilde{Q}_{1J} Q_1^J - \frac{1}{2} \tilde{Q}_{2I} Q_2^I \tilde{Q}_{2J} Q_2^J \right. \\ & \left. + \tilde{Q}_{1I} (A_1 B^1 + A_2 B^2) Q_1^I - \tilde{Q}_{2I} (B^1 A_1 + B^2 A_2) Q_2^I \right]. \end{aligned} \quad (10.3)$$

from which follow the independent F-term equations [31]

$$\begin{aligned} B^1 A_2 B^2 - B^2 A_2 B^1 + B^1 Q_1^I \tilde{Q}_{1I} - Q_2^I \tilde{Q}_{2I} B^1 &= 0, \\ B^2 A_1 B^1 - B^2 A_1 B^2 + B^2 Q_1^I \tilde{Q}_{1I} - Q_2^I \tilde{Q}_{2I} B^2 &= 0, \\ (A_1 B^1 + A_2 B^2 + Q_1^K \tilde{Q}_{1K}) Q_1^I &= 0, \\ (B^1 A_1 + B^2 A_2 + Q_2^K \tilde{Q}_{2K}) Q_2^I &= 0. \end{aligned} \quad (10.4)$$

Additional relations can be obtained by taking the transposes of eqs. (10.4).

The $\mathcal{N} = 1$ $\text{Sp}(2N) \times \text{Sp}(2N)$ gauge theory just discussed arises as the theory on a stack of D3-branes at a conifold singularity modded out by a $\mathbb{Z}_2^{\text{ori}}$ orientifold projection. In the near-horizon limit, the geometry becomes $adS_5 \times T^{11} / \mathbb{Z}_2^{\text{ori}}$. The metric on adS_5 is given by eq. (2.2); the metric on T^{11} is [38]

$$ds_{T^{11}}^2 = \frac{1}{9} (d\psi + \cos \theta_1 d\varphi_1 + \cos \theta_2 d\varphi_2)^2 + \frac{1}{6} (d\theta_1^2 + \sin^2 \theta_1 d\varphi_1^2) + \frac{1}{6} (d\theta_2^2 + \sin^2 \theta_2 d\varphi_2^2). \quad (10.5)$$

The orientifold acts by interchanging the two spheres [27, 31]: $(\varphi_1, \theta_1) \leftrightarrow (\varphi_2, \theta_2)$. The fixed point set $\theta_1 = \theta_2$, $\varphi_1 = \varphi_2$ represents the position of the O7-plane and D7-branes.

The Penrose limit

Parameterizing the angular variables in eq. (10.5) as [24, 33, 34]

$$\varphi_i = \xi - \chi_i, \quad \psi = \xi + \chi_1 + \chi_2, \quad \theta_i = \frac{\sqrt{6}}{R} y_i, \quad (10.6)$$

introducing

$$x^+ = \frac{1}{2}(t + \xi), \quad x^- = \frac{1}{2}R^2(t - \xi), \quad r = R\rho, \quad (10.7)$$

and then taking the $R \rightarrow \infty$ limit gives the standard pp-wave metric:

$$\begin{aligned} ds^2 &= R^2 \left(ds_{adS_5}^2 + ds_{T^{11}}^2 \right), \\ &\rightarrow -4dx^+ dx^- - (r^2 + y_1^2 + y_2^2)(dx^+)^2 + dr^2 + r^2 d\Omega_3^2 + dy_1^2 + y_1^2 d\chi_1^2 + dy_2^2 + y_2^2 d\chi_2^2. \end{aligned} \quad (10.8)$$

The orientifold survives the limit and acts by interchanging (y_1, χ_1) and (y_2, χ_2) . The D7-branes also survive the limit and are located at the fixed point of the orientifold, i.e. at $y_1 = y_2$, $\chi_1 = \chi_2$. This gives rise to an open-string sector of the string theory in the Penrose limit. (Contrary to the situation for the $\mathcal{N} = 2$ models, there does not appear to exist a limit in which the D7-branes disappear.)

The limiting metric and the orientifold action are identical (after a trivial coordinate redefinition) to the ones considered in ref. [8] (the gauge group on the D7-branes is different though). Thus the string theory analysis carried out in ref. [8] carries over essentially unchanged to our case, but the field theory analysis is different.

The $\mathcal{N} = 1$ $\text{Sp}(2N) \times \text{Sp}(2N)$ gauge theory has a $\text{U}(1)_R \times \text{SU}(2) \times \text{SU}(2)$ symmetry; on the string theory side, this arises from the isometry group of T^{11} . We use the following notation: J_R is the generator of $\text{U}(1)_R$, whereas J_R^3 is sum of the Cartan generators of the two $\text{SU}(2)$'s. Hence,

$$J_R^3 = -i\partial_{\varphi_1} - i\partial_{\varphi_2}, \quad J_R = -2i\partial_\psi. \quad (10.9)$$

For the Penrose limit considered above, we have $H = \Delta - J$ where $J = \frac{1}{2}J_R + J_R^3$ [24, 33, 34].

Closed string sector

The ground state in the closed string sector corresponds (in our conventions) to the $H = 0$ gauge theory operator

$$\text{tr}(B^2 A_1)^n. \quad (10.10)$$

Operators with $H = 1$ are $D_\mu(B^2 A_1)$, $B^1 A_1$, $B^2 A_2$, $(A_2)^\dagger A_1$ and $B^2(B^1)^\dagger$. (It was pointed out in ref. [24] that the conformal dimension of the latter two operators is not the naive sum of the dimensions of the constituents, but is $\Delta = 2$, resulting in $H = 1$.) In the unorientifolded case, insertions of these operators in (10.10) correspond to oscillators acting on the vacuum of the string theory [24, 33, 34].

In the orientifolded theory, the zero-momentum oscillator in the direction transverse to the orientifold plane is odd under $\mathbb{Z}_2^{\text{ori}}$, and so single oscillator states in this direction are projected out. Correspondingly, a single insertion of the operator corresponding to this direction yields a vanishing operator, as we will now show. The operators $N_{ij} = B_a^j c A_{ic}^b$ satisfy the orientifold projections [31]

$$N_{11} = J_2 N_{22}^T J_2, \quad N_{12} = -J_2 N_{12}^T J_2, \quad N_{21} = -J_2 N_{21}^T J_2, \quad (10.11)$$

by virtue of eqs. (10.1). It follows that $(N_{11} + N_{22})J_2$ is symmetric whereas $N_{12}J_2$, $N_{21}J_2$ and $(N_{11} - N_{22})J_2$ are antisymmetric. This implies that the number of insertions of $N_{11} + N_{22}$ into the vacuum state $\text{tr}(N_{12})^n$ must be even. This result corresponds to the fact that the coordinate on the string theory side corresponding to $(N_{11} + N_{22})$ changes sign under the

orientifold [31]. Similarly, the other operator which changes sign under the orientifold is $(A_2)^\dagger A_1 + B^2(B^1)^\dagger$; only an even number of insertions of this operator is possible. This follows from the fact that $[(A_2)^\dagger A_1 + B^2(B^1)^\dagger]J_2$ is symmetric due to eq. (10.1). We have thus shown that the field theory reproduces the string theory result.

Open string sector

There are three possible open string vacua formed from an $H = 0$ operator with two “quarks” at the ends:

$$Q_1^{[I} J_1(A_1 B^2)^n Q_1^{K]}, \quad Q_2^{[I} J_2(B^2 A_1)^n Q_2^{K]}, \quad Q_1^I J_1(A_1 B^2)^n A_1 Q_2^K. \quad (10.12)$$

Because of the antisymmetry of $J_2(B^2 A_1)^n$ and $J_1(A_1 B^2)^n$, the first two operators in (10.12) are antisymmetric in the interchange of I and K , and therefore belong to the adjoint **6** of $\text{SO}(4)$. This is consistent with the fact that the flavor symmetry group for the D7’s is $\text{SO}(4) \times \text{SO}(4)$, with the first two operators in (10.12) corresponding to the two independent $\text{SO}(4)$ factors (two stacks of D7-branes). The last operator in (10.12) belongs to the $(\mathbf{4}, \mathbf{4})$ representation of $\text{SO}(4) \times \text{SO}(4)$ and represents the vacuum states for strings connecting the two stacks of D7-branes.

As in the closed string sector, excitations of the open string correspond to insertions of $H = 1$ operators into the open string vacua (10.12). Operators corresponding to insertions of zero-momentum oscillators in the directions transverse to the D7-branes should be absent however [8]. As an example of this, consider the insertion of $N_{11} + N_{22}$ into $Q_2^I J_2 N_{12}^n Q_2^K$. We will now show that such an operator vanishes by using the F-term equations. From the F-term equations (10.4), it follows that $N_{11} + N_{22} + Q_2^I \tilde{Q}_{2I}$ commutes with all the N_{ij} ’s. Therefore $N_{11} + N_{22}$ can be commuted past N_{12} modulo a Q^2 term. This Q^2 term splits the operator into a “multiple trace” operator of the form $[Q_2 \cdots Q_2][Q_2 \cdots Q_2]$, which is subleading in the $1/N$ expansion. (It is natural to interpret this as a splitting of the open string into two.) Thus, $N_{11} + N_{22}$ can be moved to the end of the operator, where, again by the F-term equations (10.4), it gives zero when acting on the Q_2 (again modulo a Q^2 term, which gives a subleading contribution). Since the operator vanishes (modulo subleading terms) when using the F-term equations it is not a protected operator, consequently the corresponding zero-momentum oscillator is absent. Similarly, we expect that an insertion of $(A_2)^\dagger A_1 + B^2(B^1)^\dagger$ should give rise to an unprotected operator, reflecting the absence of the corresponding string zero-momentum state. Thus the field theory reproduces the expected string theory result [8].

11 Summary

In this paper we have studied the pp-wave limits of a variety of elliptic models with O7-planes and D7-branes, as well as the pp-wave limit of a particular orientifold of $adS_5 \times T^{11}$. The theories with D7-branes have a Penrose limit with both open and closed string sectors.

For most cases, two different Penrose limits were discussed: limit A, which gives an orbifolded pp-wave and which thus contains both an untwisted and a twisted sector; and limit B, which gives an orientifold of the pp-wave with D7-branes, and which thus contains both closed and open string sectors. In limit A (B) a remnant of the orientifold (orbifold) part

of the full orientifold group is manifested in the classification of the chiral and “near-chiral” primaries of the theories.

The two infinite series of elliptic models discussed in sections 7 and 9 allow pp-wave scaling limits analogous to the one discussed in refs. [13, 14], in which there is a compact light-cone null direction. The models that we discuss contain both open and closed strings.

In section 10 by studying a particular orientifold of $adS_5 \times T^{11}$ we were able to test the ideas of ref. [8] in a more complicated example and agreement was found between the string and field theory results.

We have not touched upon the important topic of interactions, but it would clearly be of interest to gain a better understanding of the interactions between closed and open strings and to elucidate the splitting of open strings. For some recent work on interactions in the pp-wave background and in the dual gauge theory, see e.g. refs. [22, 23, 30, 39].

Acknowledgement

HJS would like to thank the string theory group and Physics Department of Harvard University for their hospitality extended over a long period of time.

References

- [1] J. M. Maldacena, “The large N limit of superconformal field theories and supergravity.” *Adv. Theor. Math. Phys.* **2** (1998) 231–252, [hep-th/9711200](#); S. S. Gubser, I. R. Klebanov, and A. M. Polyakov, “Gauge theory correlators from non-critical string theory.” *Phys. Lett.* **B428** (1998) 105–114, [hep-th/9802109](#); E. Witten, “Anti-de Sitter space and holography.” *Adv. Theor. Math. Phys.* **2** (1998) 253–291, [hep-th/9802150](#).
- [2] D. Berenstein, J. M. Maldacena, and H. Nastase, “Strings in flat space and pp waves from $\mathcal{N} = 4$ super Yang Mills.” *JHEP* **04** (2002) 013, [hep-th/0202021](#).
- [3] R. Güven, “Plane wave limits and T-duality.” *Phys. Lett.* **B482** (2000) 255–263, [hep-th/0005061](#).
- [4] M. Blau, J. Figueroa-O’Farrill, C. Hull, and G. Papadopoulos, “A new maximally supersymmetric background of IIB superstring theory.” *JHEP* **01** (2002) 047, [hep-th/0110242](#); M. Blau, J. Figueroa-O’Farrill, C. Hull, and G. Papadopoulos, “Penrose limits and maximal supersymmetry.” *Class. Quant. Grav.* **19** (2002) L87–L95, [hep-th/0201081](#).
- [5] R. R. Metsaev, “Type IIB Green-Schwarz superstring in plane wave Ramond-Ramond background.” *Nucl. Phys.* **B625** (2002) 70–96, [hep-th/0112044](#).
- [6] R. Penrose, “Any spacetime has a plane wave as a limit.” in *Differential Geometry and Relativity*. Reidel, Dordrecht, 1976.

- [7] R. R. Metsaev and A. A. Tseytlin, “Exactly solvable model of superstring in plane wave Ramond-Ramond background.” [hep-th/0202109](#).
- [8] D. Berenstein, E. Gava, J. M. Maldacena, K. S. Narain, and H. Nastase, “Open strings on plane waves and their Yang-Mills duals.” [hep-th/0203249](#).
- [9] M. Billó and I. Pesando, “Boundary states for GS superstrings in an Hpp wave background.” [hep-th/0203028](#); C.-S. Chu and P.-M. Ho, “Noncommutative D-brane and open string in pp-wave background with B-field.” [hep-th/0203186](#); A. Dabholkar and S. Parvizi, “Dp branes in pp-wave background.” [hep-th/0203231](#); P. Lee and J.-w. Park, “Open strings in PP-wave background from defect conformal field theory.” [hep-th/0203257](#); A. Kumar, R. R. Nayak, and Sanjay, “D-brane solutions in pp-wave background.” [hep-th/0204025](#); D.-s. Bak, “Supersymmetric branes in PP wave background.” [hep-th/0204033](#); K. Skenderis and M. Taylor, “Branes in AdS and pp-wave spacetimes.” [hep-th/0204054](#); V. Balasubramanian, M.-x. Huang, T. S. Levi, and A. Naqvi, “Open strings from $\mathcal{N} = 4$ super Yang-Mills.” [hep-th/0204196](#); Y. Imamura, “Large angular momentum closed strings colliding with D-branes.” [hep-th/0204200](#); H. Takayanagi and T. Takayanagi, “Open strings in exactly solvable model of curved space-time and PP-wave limit.” [hep-th/0204234](#); P. Bain, P. Meessen, and M. Zamaklar, “Supergravity solutions for D-branes in Hpp-wave backgrounds.” [hep-th/0205106](#); M. Alishahiha and A. Kumar, “D-brane solutions from new isometries of pp-waves.” [hep-th/0205134](#); O. Bergman, M. R. Gaberdiel, and M. B. Green, “D-brane interactions in type IIB plane-wave background.” [hep-th/0205183](#); S. Seki, “D5-brane in Anti-de Sitter space and Penrose limit.” [hep-th/0205266](#); D. Mateos and S. Ng, “Penrose limits of the baryonic D5-brane.” [hep-th/0205291](#); S. S. Pal, “Solution to worldvolume action of D3 brane in pp-wave background.” [hep-th/0205303](#).
- [10] S. Gukov and A. Kapustin, “New $\mathcal{N} = 2$ superconformal field theories from M/F theory orbifolds.” Nucl. Phys. **B545** (1999) 283–308, [hep-th/9808175](#).
- [11] I. P. Ennes, C. Lozano, S. G. Naculich, and H. J. Schnitzer, “Elliptic models, type IIB orientifolds and the AdS/CFT correspondence.” Nucl. Phys. **B591** (2000) 195–226, [hep-th/0006140](#).
- [12] J. Park and A. M. Uranga, “A note on superconformal $\mathcal{N} = 2$ theories and orientifolds.” Nucl. Phys. **B542** (1999) 139–156, [hep-th/9808161](#).
- [13] S. Mukhi, M. Rangamani, and E. Verlinde, “Strings from quivers, membranes from moose.” JHEP **05** (2002) 023, [hep-th/0204147](#).
- [14] M. Alishahiha and M. M. Sheikh-Jabbari, “Strings in PP-waves and worldsheet deconstruction,” [hep-th/0204174](#).
- [15] M. Blau, J. Figueroa-O’Farrill, and G. Papadopoulos, “Penrose limits, supergravity and brane dynamics.” [hep-th/0202111](#).

- [16] E. Witten, “Baryons and branes in anti de Sitter space.” JHEP **07** (1998) 006, [hep-th/9805112](#).
- [17] E. Floratos and A. Kehagias, “Penrose limits of orbifolds and orientifolds.” [hep-th/0203134](#).
- [18] A. Sen, “F-theory and Orientifolds.” Nucl. Phys. **B475** (1996) 562–578, [hep-th/9605150](#); T. Banks, M. R. Douglas, and N. Seiberg, “Probing F-theory with branes.” Phys. Lett. **B387** (1996) 278–281, [hep-th/9605199](#); O. Aharony, J. Sonnenschein, S. Yankielowicz, and S. Theisen, “Field theory questions for string theory answers.” Nucl. Phys. **B493** (1997) 177–197, [hep-th/9611222](#); M. R. Douglas, D. A. Lowe, and J. H. Schwarz, “Probing F-theory with multiple branes.” Phys. Lett. **B394** (1997) 297–301, [hep-th/9612062](#).
- [19] A. Fayyazuddin and M. Spalinski, “Large N superconformal gauge theories and supergravity orientifolds.” Nucl. Phys. **B535** (1998) 219–232, [hep-th/9805096](#); O. Aharony, A. Fayyazuddin, and J. M. Maldacena, “The large N limit of $\mathcal{N} = 2, 1$ field theories from three- branes in F-theory.” JHEP **07** (1998) 013, [hep-th/9806159](#).
- [20] N.-w. Kim, A. Pankiewicz, S.-J. Rey, and S. Theisen, “Superstring on pp-wave orbifold from large- N quiver gauge theory.” [hep-th/0203080](#).
- [21] M. R. Douglas and G. W. Moore, “D-branes, Quivers, and ALE Instantons.” [hep-th/9603167](#).
- [22] C. Kristjansen, J. Plefka, G. W. Semenoff, and M. Staudacher, “A new double-scaling limit of $\mathcal{N} = 4$ super Yang-Mills theory and PP-wave strings.” [hep-th/0205033](#).
- [23] N. R. Constable *et. al.*, “PP-wave string interactions from perturbative Yang-Mills theory.” [hep-th/0205089](#).
- [24] N. Itzhaki, I. R. Klebanov, and S. Mukhi, “PP wave limit and enhanced supersymmetry in gauge theories.” JHEP **03** (2002) 048, [hep-th/0202153](#).
- [25] M. Alishahiha and M. M. Sheikh-Jabbari, “The PP-wave limits of orbifolded $AdS_5 \times S^5$.” [hep-th/0203018](#).
- [26] T. Takayanagi and S. Terashima, “Strings on orbifolded pp-waves.” [hep-th/0203093](#).
- [27] S. G. Naculich, H. J. Schnitzer, and N. Wyllard, “ $1/N$ corrections to anomalies and the AdS/CFT correspondence for orientifolded $\mathcal{N} = 2$ orbifold models and $\mathcal{N} = 1$ conifold models.” [hep-th/0106020](#).
- [28] L. Susskind, “Another conjecture about M(atrix) theory.” [hep-th/9704080](#).
- [29] M. Cvetič, H. Lü, and C. N. Pope, “Penrose limits, pp-waves and deformed M2-branes.” [hep-th/0203082](#); J. Michelson, “(Twisted) toroidal compactification of pp-waves.” [hep-th/0203140](#).

- [30] R. Gopakumar, “String interactions in PP-waves.” [hep-th/0205174](#).
- [31] S. G. Naculich, H. J. Schnitzer, and N. Wyllard, “A cascading $\mathcal{N} = 1$ $\text{Sp}(2N+2M) \times \text{Sp}(2N)$ gauge theory.” [hep-th/0204023](#).
- [32] I. R. Klebanov and E. Witten, “Superconformal field theory on threebranes at a Calabi-Yau singularity.” Nucl. Phys. **B536** (1998) 199–218, [hep-th/9807080](#).
- [33] J. Gomis and H. Ooguri, “Penrose limit of $\mathcal{N} = 1$ gauge theories.” [hep-th/0202157](#).
- [34] L. A. Pando Zayas and J. Sonnenschein, “On Penrose limits and gauge theories.” JHEP **05** (2002) 010, [hep-th/0202186](#).
- [35] K. Oh and R. Tatar, “Orbifolds, Penrose limits and supersymmetry enhancement.” [hep-th/0205067](#).
- [36] F. Bigazzi, A. L. Cotrone, L. Girardello, and A. Zaffaroni, “pp-wave and non-supersymmetric gauge theory.” [hep-th/0205296](#).
- [37] H. J. Schnitzer and N. Wyllard, “An orientifold of $adS_5 \times T^{11}$ with D7-branes, the associated α'^2 -corrections, and their role in the dual $\mathcal{N} = 1$ $\text{Sp}(2N + 2M) \times \text{Sp}(2N)$ gauge theory.” [hep-th/0206071](#).
- [38] P. Candelas and X. C. de la Ossa, “Comments on conifolds.” Nucl. Phys. **B342** (1990) 246–268.
- [39] M. Spradlin and A. Volovich, “Superstring interactions in a pp-wave background.” [hep-th/0204146](#); A. Parnachev and D. A. Sahakyan, “Penrose limit and string quantization in $AdS_3 \times S^3$.” [hep-th/0205015](#); D. J. Gross, A. Mikhailov, and R. Roiban, “Operators with large R charge in $\mathcal{N} = 4$ Yang-Mills theory.” [hep-th/0205066](#); C.-S. Chu, P.-M. Ho, and F.-L. Lin, “Cubic string field theory in pp-wave background and background independent Moyal structure.” [hep-th/0205218](#); Y.-j. Kiem, Y.-b. Kim, S.-m. Lee, and J.-m. Park, “pp-wave/Yang-Mills correspondence: An explicit check.” [hep-th/0205279](#); M.-x. Huang, “Three point functions of $\mathcal{N} = 4$ Super Yang Mills from light cone string field theory in pp-wave.” [hep-th/0205311](#); C.-S. Chu, V. V. Khoze, and G. Travaglini, “Three-point functions in $\mathcal{N} = 4$ Yang-Mills theory and pp-waves.” [hep-th/0206005](#).